

THURROCK

(EX)

THPE94
328/95

Union Railways Limited
Channel Tunnel Rail Link

Purfleet, Thurrock, Essex

Project Code: ARCTHPE 94

Archaeological Evaluation Report

TIS No. 192/84-10411

OXFORD ARCHAEOLOGICAL UNIT

February 1996

Union Railways Limited
Channel Tunnel Rail Link

PURFLEET, THURROCK, ESSEX
PROJECT CODE: ARCTHPE 94
ARCHAEOLOGICAL EVALUATION REPORT

TIS No. 192/84 - 10411

| |
|-----------------------|
| Prepared by: Date: |
| Checked by: Date: |
| Approved by: Date: |

OXFORD ARCHAEOLOGICAL UNIT

February, 1996

PURFLEET, THURROCK, ESSEX
ARCHAEOLOGICAL EVALUATION

LIST OF CONTENTS

| | | |
|------|---|----|
| | SUMMARY | 1 |
| 1 | BACKGROUND | 1 |
| 1.1 | Introduction | 1 |
| 1.2 | Reason for the Project | 1 |
| 1.3 | Construction Work on the Purfleet site | 3 |
| 1.4 | Geology, Landscape and Landuse | 3 |
| 1.5 | Archaeological Background | 4 |
| 2 | AIMS | 6 |
| 3 | METHODS | 7 |
| 3.1 | General | 7 |
| 3.2 | Borehole Investigation | 7 |
| 3.3 | Test Pit Investigation | 9 |
| 3.4 | Survey | 14 |
| 4 | RESULTS: BOREHOLE INVESTIGATION | 14 |
| 4.1 | General | 14 |
| 4.2 | ARC0006 SA | 16 |
| 4.3 | ARC0007 SA | 19 |
| 4.4 | ARC0008 SA | 22 |
| 4.5 | ARC0009 SA | 25 |
| 4.6 | ARC0010 SA | 30 |
| 4.7 | ARC0011 SA | 33 |
| 4.8 | ARC0012 SA | 37 |
| 4.9 | ARC0013 SA | 41 |
| 4.10 | ARC0014 SA | 44 |
| 4.11 | ARC0015 SA | 49 |
| 5 | LOCATION OF TEST PITS | 52 |
| 6 | RESULTS: TEST PIT INVESTIGATION | 53 |
| 6.1 | ARC2001 TP | 53 |
| 6.2 | ARC2002 TP | 57 |
| 6.3 | ARC2003 TP | 60 |
| 6.4 | ARC2004 TP | 63 |
| 6.5 | Summary of Artefacts and Samples Recovered | 66 |
| 7 | SPECIALIST REPORTS | 68 |
| 7.1 | Flint Report | 68 |
| 7.2 | Pollen Analysis | 72 |
| 7.3 | Prehistoric Pottery | 74 |
| 7.4 | Molluscs, Charred Plant Remains and Bones | 75 |
| 8 | SUMMARY OF RESULTS | 76 |
| 8.1 | General Stratigraphic Sequence | 76 |
| 8.2 | Test Pits | 76 |
| 8.3 | Biological and Artefactual Material | 78 |
| 9 | CONCLUSIONS | 80 |
| 9.1 | Pleistocene | 80 |
| 9.2 | Later Prehistoric | 81 |
| 10 | SIGNIFICANCE OF DEPOSITS | 83 |
| 10.1 | General Considerations | 83 |
| 10.2 | The Purfleet Pleistocene and Palaeolithic Complex | 84 |
| 11 | BIBLIOGRAPHY | 88 |

APPENDIX

| | |
|---------------------------------------|----|
| Sample Statistics for Boreholes | 89 |
|---------------------------------------|----|

List of Tables

| | |
|----------|--|
| Table 1 | Summary of the key stratigraphic elements that were expected to be located during test-pitting, with depths to critical Pleistocene sediments and estimated Palaeolithic archaeological potential. |
| Table 2 | Deposits recorded in ARC2001 TP |
| Table 3 | Deposits recorded in ARC2002 TP |
| Table 4 | Deposits recorded in ARC2003 TP |
| Table 5 | Deposits recorded in ARC2004 TP |
| Table 6 | Summary of excavated spits, artefact recovery, and bulk samples |
| Table 7 | Summary of flint collection |
| Table 8 | Recovered flint collection divided according to individual spits |
| Table 9 | Pollen data obtained from 32-33cm (ARC2001 TP, monolith 8) |
| Table 10 | Assessment of the Significance of the Deposits |

List of Figures

| | |
|---------|--|
| Fig. 1 | Site location map |
| Fig. 2 | Test Pit and Borehole Location Plan |
| Fig. 3 | Symbols used in stratigraphic record sheet logging |
| Fig. 4 | Stratigraphic record sheet for Borehole ARC0006 SA |
| Fig. 5 | Stratigraphic record sheet for Borehole ARC0007 SA |
| Fig. 6 | Stratigraphic record sheet for Borehole ARC0008 SA |
| Fig. 7 | Stratigraphic record sheet for Borehole ARC0009 SA |
| Fig. 8 | Stratigraphic record sheet for Borehole ARC0010 SA |
| Fig. 9 | Stratigraphic record sheet for Borehole ARC0011 SA |
| Fig. 10 | Stratigraphic record sheet for Borehole ARC0012 SA |
| Fig. 11 | Stratigraphic record sheet for Borehole ARC0013 SA |
| Fig. 12 | Stratigraphic record sheet for Borehole ARC0014 SA |
| Fig. 13 | Stratigraphic record sheet for Borehole ARC0015 SA |
| Fig. 14 | South-west facing section of Test Pit ARC2001 TP |
| Fig. 15 | South-west facing section of Test Pit ARC2002 TP |
| Fig. 16 | South-west facing section of Test Pit ARC2003 TP |
| Fig. 17 | South-west facing section of Test Pit ARC2004 TP |
| Fig. 18 | Summarised stratigraphy along the route corridor based on selected boreholes |
| Fig. 19 | Summarised stratigraphy across southern edge of palaeochannel |

List of Plates

| | |
|----------|---|
| Plate 1 | The Strata Investigation drilling team, sub-contracted by the GSF for work at Purfleet, drilling at borehole ARC 0012 SA (2/8/95). |
| Plate 2 | Stratigraphic logging of split U4/U100 core samples in the Wolfson Archaeological Science Laboratories, University College London |
| Plate 3 | Excavation of Test Pit ARC2003 TP |
| Plate 4 | U4 4 core at 2.20-2.65m below ground surface, borehole ARC0009 SA showing laminated sands and silts overlying a thin gravel bed. |
| Plate 5 | U4 3 core at 1.60-2.05m below ground surface, borehole ARC 0009 SA showing finely laminated sands and silts. |
| Plate 6 | Borehole ARC 0011 SA, U4 3, 22-23cm within U4/U100 core, smooth surface quartz grain (magnification x250). |
| Plate 7 | Borehole ARC 0011 SA, U4 4, 21-22cm within U4/U100 core, clay-coated quartz grains (magnification x170). |
| Plate 8 | Unsieved bulk sample B4, ARC 0012 SA, 1.80-2.00m depth. |
| Plate 9 | U4 2 core at 1.10-1.55m below ground surface (lower part), borehole ARC0014 SA, showing sharp contact between the sands and underlying chalk bedrock. |
| Plate 10 | U4 2 core 1.10-1.55m below ground surface (upper part), borehole ARC0014 SA, showing laminated sands and silts. |
| Plate 11 | Unsieved bulk sample B3, ARC 0015 SA, 2.10-2.40m depth. |
| Plate 12 | Bifacial hand-axes (top) and biface manufacturing flakes (bottom) from ARC2004 TP |

PURFLEET, THURROCK, ESSEX
TQ 562784

ARCHAEOLOGICAL EVALUATION

SUMMARY

As part of a larger programme of archaeological investigation along the route of the Channel Tunnel Rail Link, Union Railways Ltd commissioned the Oxford Archaeological Unit to undertake a field evaluation of 0.5ha of land immediately to the south of the Purfleet By-pass, Purfleet, Thurrock, Essex in July to September 1995 (Project Code ARCTHPE 94). Purfleet is of national, if not international, importance for Palaeolithic archaeology and pleistocene geology, as recognised from previous investigations of chalk and gravel pits in the locality. Quarrying has significantly depleted this resource. The deposits of interest represent the fill of an ancient river channel, aligned east to west, which is crossed obliquely by the CTRL route at a point where the railway runs in a cutting. In this evaluation, the sedimentary stratigraphy of the deposits has been recorded from a series of ten boreholes and four test pits. A considerable number of Palaeolithic flint artefacts were recovered, largely from a slope wash or solifluction deposit eroded from a chalk cliff forming the southern edge of a palaeochannel of the Thames. Lithic material was also recovered from fluvial sediments at the margin of the palaeochannel. In both cases the flint was in relatively good condition, indicating that it had undergone little post-depositional re-working. Pollen was recovered from fine-grained sediments, but molluscs and bones were all but absent from the deposits investigated.

In addition, some evidence of later prehistoric occupation was recorded, although the nature of the evaluation precluded detailed interpretation of the features recorded.

1 BACKGROUND

1.1 Introduction

1.1.1 The Oxford Archaeological Unit undertook a two stage archaeological evaluation between 24th July and 4th August 1995 and between the 18th and 22nd September 1995, on behalf of Union Railways Ltd (URL) on land at the former Esso oil depot immediately to the south of the Purfleet By-pass, Purfleet, Thurrock, Essex (Figure 1). The evaluation forms part of a programme of archaeological investigation along the line of the Channel Tunnel Rail Link, the aim of which is to appraise the impact of the construction of the new railway upon the cultural heritage.

1.1.2 The work was carried out in accordance with a Written Scheme of Investigation (WSI), detailing the scope and method of the evaluation. Pleistocene deposits within the Purfleet area have been extensively quarried, allowing scientific and archaeological investigations to be carried out on them. The southern edge of a palaeochannel of the Thames crosses the evaluation site, and previous studies have shown that Palaeolithic artefacts are most densely distributed towards the southern edge of the pleistocene deposits.

1.2 Reason for the Project

1.2.1 HM Government has determined that a new railway should be built to connect London mainline railway stations and the Channel Tunnel. The project involves extensive construction work, including cuttings, tunnels and embankments.




Figure 1. Site location plan showing distribution of Pleistocene sand and gravels and the transect line of boreholes drilled during the GSF site investigation.

- 1.2.2 An Environmental Statement has been prepared. This examines the impact of the project on the natural and built environment. The OAU has provided detailed archaeological input to this document (Assessment of Historic and Cultural Effects, Vols 1-4, URL 1994) and a number of sites have been identified where the proposed route of the rail link will affect areas of particular known or potential archaeological sensitivity. Archaeological evaluation of these areas is necessary so that detailed mitigation strategies can be devised for the archaeology (references to these features in the Purfleet area are given here as OAU No.... and may be found in Volume 1 Route Corridor; Volume 2 Maps; Volume 3 Gazetteer; and Volume 4 Appendix B).
- 1.2.3 The CTRL Assessment of Historical and Cultural Effects (URL 1994) identifies the Pleistocene channel deposits at Purfleet, with the co-occurrence of three distinct artefactual horizons in association with deposits containing molluscan, faunal and pollen evidence, as of at least national importance.
- 1.3 Construction Work on the Purfleet Site**
- 1.3.1 The route crosses the Borough of Thurrock to approach the north portal of a new tunnel under the river Thames. Construction will involve cuttings, embankments and ground-level works. Some of the route through the area north of Purfleet crosses ground which has been subject to mineral extraction in the past, and is therefore of little or no archaeological sensitivity. The evaluation area, however, contains a significant extent of substantially intact land (although part of the site appears to have been affected by extraction). The railway will run through the site in cutting up to 8 m deep at the south-east end.
- 1.4 Geology, Landscape and Landuse**
- 1.4.1 The Aveley Marshes end at Purfleet with the commencement of the chalk outcrop that extends through most of the ancient parish of West Thurrock. The Mar Dyke follows the line of a former, more sinuous course of the Thames. Extensive overlying gravels and other deposits on both the north and south sides of the Dyke provide evidence of the extent of the former valley. The CTRL affects the deposits on the south side.
- 1.4.2 The fluvial deposits at Purfleet are part of the Corbets Tey Gravel, the middle of three gravel terraces which have been recognised in the Lower Thames.
- 1.4.3 Elements of a composite stratigraphic sequence of pleistocene deposits determined by previous workers were expected in the area of study. The composite profile constructed by Bridgland (1994) and based on Hollin (1977) shows 5 major units are present above Chalk bedrock in the area. It is thought likely that these units will inter-digitate and change laterally. This is because the gravel and sand deposits exhibit bedding structures and sedimentological trends consistent with deposition in fluvial channels which are subject to rapid channel shifts, variable flow regimes and rapid shifts in the loci of sedimentation. The accepted model for the development of the Thames fluvial sequences (Bridgland 1994, 17-19) indicates that coarse sands and gravels are typically laid down in cold climate conditions while the finer

grained interbedded sands, silts and clays overlying the gravels indicate temperate floodplain environments.

- 1.4.4 The analysis of oxygen isotopes in fossil plankton in deep ocean cores provides an indication of ice-cap size and therefore of climate. Warm and cold episodes are numbered backwards from the present interglacial - Oxygen Isotope Stage 1. The oldest of the Lower Thames interglacials, the Hoxnian, is correlated with Stage 11. Bridgland (1994, 13) has correlated the temperate deposits within the Corbets Tey Gravel to Oxygen Isotope Stage 9.
- 1.4.5 The Pleistocene sediments are banked against a chalk cliff line that exists towards the eastern end of the evaluation area.
- 1.4.6 Part of the north-western end of the site has been designated as a Site of Special Scientific Interest (SSSI), known as Purfleet Chalk Pits, because of its potential to demonstrate the relationship of the Purfleet deposits to other significant deposits within the Thames terrace sequence (Bridgland 1994).
- 1.4.7 The evaluation area mainly consists of rough pasture, scrub and a few mature trees, with an area of hard-standing towards the centre.
- 1.5 **Archaeological background**
- 1.5.1 The Purfleet area is known to contain archaeological remains from the Palaeolithic up to the Roman periods.
- 1.5.2 **Palaeolithic**
- 1.5.2.1 The Palaeolithic evidence consists of flint artefacts and palaeo-environmental remains recovered from Pleistocene geological deposits. These deposits consist of a sequence of stratified clays, silts, sands and gravels lying in a channel cut into the north-facing chalk slope to the south of the Mar Dyke. This channel and its infilling Pleistocene deposits represent an early course of the Thames. A substantial part of these deposits has been removed by quarrying.
- 1.5.2.2 Palaeolithic flint artefacts were collected by Snelling from Botany Pit (TQ 556785) in 1961 (URL 1994, OAU No. 1521). This flint industry has been attributed as "Proto-Levalloisian" and dates to between 200,000 and 300,000 years ago. Snelling recovered this assemblage from a deposit of gravel approx. 3.5 m thick resting on chalk bedrock at approx. 12 m OD (Wymer 1968, 312-313).
- 1.5.2.3 More detailed investigations of the Mar Dyke channel deposits (OAU No. 1516) were carried out less than 1 km to the east by Palmer in the late 1960s at Greenlands Quarry (TQ 566785) and Bluelands Quarry (TQ 570787). Palmer (1975, 1-13) recovered Palaeolithic artefacts from channel deposits in both quarries. These artefacts occurred in three distinct bands of gravel within the Pleistocene sequence. The artefacts from each gravel band have been attributed to, respectively: the Levalloisian industry (Gravel 1), the Clactonian and Acheulian industries (Gravel 2) and the Clactonian industry (Gravel 3).

Palmer identified these gravel-bands within a deep sequence of Pleistocene deposits filling the middle of the Mar Dyke channel. At this point the base of the channel was cut into chalk bedrock at 6.7 m OD, and the top of the Pleistocene sequence occurred at 13.75 m OD. In between the gravel-bands 3 and 2, between 2 and 1, and over gravel-band 1, are complex sequences of clays, silts and sands. The lower clays and silts (between gravel-bands 3 and 2) have been shown to contain both mollusc shells and pollen grains. The presence of molluscan remains suggests suitable conditions for the preservation of faunal remains, and in particular small mammals (voles, shrews and mice).

- 1.5.2.4 A recent re-examination of the Botany and Esso pits by David Bridgland as part of a geological study of the CTRL (Bridgland 1993) suggests that Palaeolithic artefacts are most strongly distributed towards the southern edge of the Pleistocene deposits where these are banked against the chalk anticline.
- 1.5.3 Mesolithic and Neolithic
- 1.5.3.1 Archaeological remains from these periods have been discovered at several locations in the area. At the west end of Greenlands Quarry (TQ 56407850) pottery (possibly Neolithic), flint artefacts (Mesolithic/Neolithic) and two possible post-holes (unattributed) were found in the top metre of the deposits exposed in the quarry section (OAU 1518). A prolific Mesolithic flint industry (OAU 1517) was also recovered from the topmost part of the section at the northeast corner of Greenlands Quarry (TQ 56857853). An assemblage of over 300 flint artefacts (OAU No. 1961), including flakes and a polished axe of likely Mesolithic and/or Neolithic age, was recovered from the topsoil at Beacon Hill (TQ 557782) during an archaeological salvage operation after the destruction of the hill-top by quarrying in 1969 (Caldwell 1971, 58).
- 1.5.4 Bronze Age
- 1.5.4.1 The only unambiguous location of Bronze Age archaeological remains is at the top of Beacon Hill (TQ 557782), where a Bronze Age cinerary urn (OAU No. 1961) was recovered during the 1969 archaeological salvage operation (Caldwell 1971, 58).
- 1.5.5 Iron Age and Roman
- 1.5.5.1 Isolated finds of pottery, bricks, tiles and loomweights attributed to the Roman and/or Iron Age periods (OAU No. 1522) have regularly been found in the upper parts of the old quarry faces around Botany Pit (TQ 556785). An inspection of "the old working face" in 1956 (Gilman 1956, 12-14) showed traces of pits and ditches containing pottery, bones and much charcoal (OAU No. 1523). These features can probably be linked with the isolated Roman/Iron Age finds common in the Botany Pit. No precise locations have been recorded for these archaeological remains, although they most likely came from the high west-facing side of the quarry.

- 2 **AIMS**
- 2.1 The Written Scheme of Investigation (URL 1995c) outlined thirteen principal aims for the evaluation.
- 2.2 The evaluation aims were as follows:
- 2.2.1 To assess the sedimentological character, and horizontal and vertical extent, of affected Pleistocene/Palaeolithic deposits.
- 2.2.2 To assess the vertical and lateral distribution of palaeolithic artefacts within the Pleistocene levels.
- 2.2.3 To examine the Pleistocene/Palaeolithic levels to the depth of impact from construction works by means of boreholes and by test pits.
- 2.2.4 To assess the potential of the Pleistocene/Palaeolithic and other deposits for artefactual and environmental remains.
- 2.2.5 To assess the relationship of Palaeolithic archaeological deposits, including artefacts and ecofacts, to the Pleistocene lithostratigraphic and sedimentary sequence.
- 2.2.6 To determine the potential for post-Palaeolithic archaeological deposits and, if such deposits are defined, to assess their date, character, extent, quality and condition.
- 2.2.7 To relate all archaeological deposits found to other discoveries in the locality.
- 2.2.8 To critically review the local, regional, national and (where relevant) international significance of such archaeological deposits as are revealed.
- 2.2.9 To contribute towards proposals for mitigation of impact on such archaeological deposits as are revealed and/or can be predicted from the evaluation evidence.
- 2.2.10 To make a full graphic, photographic and written record of the evaluation.
- 2.2.11 To communicate the results of the evaluation to the client (and through them to the statutory consultees) in the form of a suitably illustrated report which shall be lodged with the County Sites and Monuments Record within one year of the end of fieldwork.
- 2.2.12 To prepare an archive of the evaluation project, to be deposited in an approved museum within a timescale to be agreed with the County Archaeologist, taking due account of the potential for further fieldwork.
- 2.2.13 To deposit the finds with the archive (subject to the agreement of the landowner and, where relevant, any decisions under Treasure Trove law).

3 METHODS

3.1 General

3.1.1 A detailed statement of the methods used in the evaluation is contained within Section 3 of the Written Scheme of Investigation (URL 1995c), the accompanying Methods Statement (URL 1995a) and the Site Safety Plan and COSHH Statement (URL 1995b). The following is only intended to amplify certain aspects of the evaluation methodology.

3.1.2 In order to evaluate the thick and deeply buried pleistocene deposits, a staged approach to the evaluation was devised

3.1.3 The first stage of work took the form of a borehole investigation, carried out by the Geoarchaeological Service Facility (GSF) of the Institute of Archaeology, University College, London at the request of the Oxford Archaeological Unit. A series of shell and auger core samples (see Figure 2) were taken in order to provide an assessment of the general profile and character of the lithographic and sedimentary depositional sequence and to inform the location of test-pits during the second stage of evaluation.

3.1.4 The second stage of work involved the excavation of a series of test-pits (see Figure 2), primarily for the recovery of artefacts and ecofacts, and to provide an opportunity for examination of the depositional sequence *in situ*.


3.2 Borehole investigation

3.2.1 A shell and auger percussion drill rig, capable of drilling and casing to depths of c.20m below ground surface through a variety of sediment types including sands and gravels, was used to drill an array of 10 boreholes along the proposed CTRL route corridor (Figure 2). This was carried out by Strata Investigation under sub-contract to GSF (see Plate 1).

3.2.2 In all cases, irrespective of apparent depth of Pleistocene sediments, boreholes were drilled to 10.0m depth from the surface to ensure that bedrock had been penetrated. As Pleistocene sediment thickness was typically shallower than anticipated, the Chalk was drilled to depths in excess of 5.5m in all cases.

3.2.3 Individual boreholes were recorded in detail during drilling. Where possible U4/U100 cores were recovered. Recovery quality varied both down profile and across the site. Dense ground within the gravels precluded the use of U4/U100 cores in certain parts of the sequence. Elsewhere the extremely unconsolidated nature of the sands resulted in U4/U100 cores falling apart on extraction from the ground. Where sampling using U4/U100 cores was not possible open shelling of the hole was undertaken and sediments logged and taken as bulk samples.

3.2.4 All boreholes were logged in accordance with geological descriptive standards commonly in use by the GSF staff. These also accord with, or extend, the standards used by the Museum of London Archaeological Service.



scale 1:5000

Figure 2. Test Pit and Borehole Location Plan.

- 3.2.5 Where possible cores were extruded using an hydraulic core extruder. Where sample cores were overcompacted due to the nature of the sediment sampled (e.g. U4₂ 1.10-1.55m, borehole ARC 0014 SA), cores were removed by cutting through the core tubes longitudinally.
- 3.2.6 Extruded cores were split and individual halves retained in split plastic piping or the split U4/U100's. Both core halves were cleaned carefully using a knife or scalpel and the cleaned faces examined (see Plate 2). One half of the core was selected for detailed study and photography (see Plates 4, 5, 8 and 9).
- 3.2.7 Individual cores were subsequently described in detail using standard sedimentological terminology to describe colour, composition, bedding and grain sizes. In addition other features, such as the presence of clay coatings on clasts and sand grains, zones of reddening, blocky structure to the sediment and trace fossils indicative of plant rooting/bioturbation (possibly indicating pedogenic activity and/or the presence of a buried landsurface) were actively sought.
- 3.2.8 As described above at Section 3.2.7, considerable emphasis was placed on the recognition of i) incipient pedostratigraphic features and ii) stratigraphies containing bioassemblages. When it became clear at an early stage in the bulk processing that faunal material (in particular molluscs) were not present within the cores, extra attention and resources were allocated to other possible avenues of identifying incipient pedogenesis. For this reason a previously untried route to this objective - using reddened clay coatings on quartz sand grains as a criterion identifiable by Scanning Electron Microscopy (SEM) - was deployed on a small number of selected samples from borehole ARC 0011 SA (see Plates 6 and 7).
- 3.2.9 All cores were photographed using Pentax ILX SLR cameras with 50mm SMC Pentax-A Macro 1:2.8 lenses. The colour slide film (Kodak Ektachrome 64T, 64 ASA) was tungsten balanced, while the colour print film was used with a 80A Colour correction filter to produce tungsten balancing. An International Federation of Rock Art Organisations (IFRAO) colour standard chart, with standard 10cm scale, was inserted in each frame to standardise the colour photography.
- 3.3 The Test Pit Investigation**
- 3.3.1 Four test-pits were excavated using a mechanical excavator equipped with a 0.9m wide ditching bucket (see Plate 3). All the test pits were 4m long and 1.7m wide in order to accommodate a trench-box shoring system once a depth of 1.2m had been reached. The trench-box was lowered in stages as each test pit was excavated. This technique ensured that the short end-sections of each test pit were open to observation. None of the test pits exceeded 2.95m in depth.
- 3.3.2 A resident engineer was on site at all times to advise on and monitor the safety aspects of the excavation. Test pit ARC2002aTP was resited after made ground was encountered in its initial position, and the final test pit, ARC2002TP, was excavated through unconsolidated sediments. A partial



Plate 1. The Strata Investigation drilling team, sub-contracted by the GSF for work at Purfleet, drilling at borehole ARC 0012 SA (2/8/95).



Plate 2. Stratigraphic logging of split U4/U100 core samples in the Wolfson Archaeological Science Laboratories, University College London.

collapse occurred at a depth of 2.4m below ground surface and the test pit was abandoned and backfilled.

- 3.3.3 The location of test pits ARC2001TP, ARC2002TP, and ARC2003TP was determined in advance on the basis of the results of the coring operation (Figure 2). The location of ARC2004TP was determined on site after an initial review of the deposits recorded in ARC2001TP.
- 3.3.4 The test pits were excavated in spits, nominally of 0.25m depth. The spoil from each spit was stored separately to allow hand-sorting and sieving of the material for the recovery of artefacts. The topsoil was stored separately and an additional spoil heap was required for stratigraphically mixed material. In practice, the spit depth was generally greater than 0.25m due to the need to ensure a clean surface before the next spit was excavated. The excess spoil generated in this way was placed separately on a waste heap. Each spit was numbered separately.
- 3.3.5 Sediments were recorded by M Bates and C Pine of GSF in accordance with geological descriptive standards commonly in use by the GSF staff. These also accord with, or extend, the standards used by the Museum of London Archaeological Service.
- 3.3.6 One section in each test-pit was drawn at a scale of 1:20. In test pits ARC2001TP, ARC2002TP, and ARC2003TP, the end section of the test pit was felt to be representative of the deposits as a whole; in ARC2004TP, with more complex stratigraphy, the side face was drawn as a cumulative section as the trench-box was gradually lowered.
- 3.3.7 Approximately 100 litres of spoil from each spit was coarse sieved through 10mm and 4mm meshes. In addition, a variable proportion (generally about 300 litres) of the spoil was hand-sorted. M White of Cambridge University was present on site to supervise the recovery of the flint artefactual material and has produced a specialist assessment of the material recovered (Section 7.1).
- 3.3.8 Bulk samples for environmental analysis and recovery of small artefacts were taken from each major stratigraphic unit. Forty litre samples were taken from *in situ* deposits where possible, or from the excavated spoil where safety considerations dictated. A ten litre sub-sample of each was processed by mechanical flotation to assess the potential for recovery of molluscs, bone and small artefacts such as flint chips. The non-floating residue was sieved through 10mm, 4mm and 0.5mm meshes and sorted for artefacts and bone. The flots were scanned under a binocular microscope by Dr M Robinson and the results are given in Section 7.4.
- 3.3.9 Column samples were taken from deposits to assess the potential for preservation of pollen, diatoms, and ostracods (monolith samples), as well as for micromorphological analysis (kubiena samples). Two of the monolith samples have been assessed for pollen preservation and the results are given in Section 7.2.



Plate 3. Excavation of Test Pit ARC 2003 TP.

3.4 Survey

- 3.4.1 The borehole and test pit locations were surveyed by Simmons Survey Partnership for URL. This data was supplied to the OAU.

4 RESULTS: BOREHOLE INVESTIGATION

4.1 General

- 4.1.1 The results for each borehole in turn are presented as a description of the stratigraphy: an assessment of geoarchaeological potential; and a summary of the results.
- 4.1.2 The integrated lithostratigraphic log is displayed as a stratigraphic column of constant width and scaled to a common depth scale (see Figure 3 for symbols used and Figures 4 to 13 for individual core descriptions). A second column adjacent to the lithological column indicates trends in grain sizes where increased coarseness of the sediment is represented by an increase from left to right across the horizontal axis. This column also portrays other stratigraphic data such as structural bedding features, contact types etc. A subsequent column logs the particle size of the largest clasts recorded within the gravel size fraction (an absence of record in this column indicates that no gravel sized clasts were present in the stratigraphy at this point). This form of data portrayal is useful as it rapidly indicates trends inherent in the stratigraphy and enables comparison with other sequences.
- 4.1.3 Further columns indicate the presence of rooting and bioturbation, the presence of clay coatings on the clasts and sand grains and the presence of biostratigraphic information (e.g. molluscs). The final columns contain information on facies interpretation and archaeological potential.
- 4.1.4 Figures 4 to 13 have been drawn to concentrate on the unconsolidated sediment above the Chalk bedrock (even though each borehole was drilled to a depth of at least 10.0m below ground surface). U4/U100 and bulk samples taken from within the Chalk bedrock are therefore not shown on the stratigraphic record sheets. The total number of U4/U100 core and bulk samples (from the unconsolidated sediments and from the Chalk), for each borehole, is given in the text (see also Appendix 1). The U4 and bulk samples are referred to in the form U4₁, U4₂, U4₃; B₁, B₂, B₃ etc, for each borehole.






Figure 3. Symbols used in stratigraphic record sheet logging.

4.2 ARC 0006 SA (Figure 4)

4.2.1 Stratigraphy

4.2.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.0m below ground surface. Ground level at the top of the borehole was +18.812m O.D. and the base of the borehole was at +8.812m O.D. A total depth of c. 2.08m of unconsolidated sediments was found overlying rotted Chalk bedrock.

4.2.1.2 One U4/U100 core sample was taken and 8 bulk samples were recovered. Location of samples retrieved from the unconsolidated sediments relative to the stratigraphy is shown in Figure 4.

4.2.1.3 The Chalk bedrock lies at a depth of c. 2.08m (+16.732m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment is sharp and was clearly seen during logging of the borehole in the field.

4.2.1.4 Immediately above the Chalk bedrock is the first of the three groups of sediments present within the borehole. The first is a thin bed (3.0cm) of coarse/medium sand with small flint fragments.

4.2.1.5 The second group of sediments present within the borehole is formed by a sequence of sands between core depths of 2.05m and 1.00m (+16.762m O.D. and +17.812m O.D.). The group is dominated by silty sand units with angular flint and sub-angular chalk gravel clasts up to a maximum of 5.0cm mean diameter.

4.2.1.6 The third of the groups present within the borehole is formed by silty sands, sands and a chalky gravel unit and is present between core depths of 1.00m (+17.812m O.D.) and the base of the fill at 0.30m (+18.512m O.D.). The upper part of this group of sediments contains charcoal and within the unit immediately below this there are probable infilled root channels between 0.60 and 0.40m (+18.212m and +18.412 O.D.).

4.2.1.7 The sediments within sample U4₁ (0.65-0.40m below ground surface, +18.162 to +18.412m O.D.) contained some CaCO₃ (reacted with dilute hydrochloric acid, see Appendix V). It is currently unknown whether CaCO₃ is present within the sediment below U4₁.

4.2.2 Interpretation

4.2.2.1 The lowest group of sediments (sand found directly over the Chalk), was thin and only observed during the field logging. It is possible that this deposit was laid down by running water. The second group of sediments present within the borehole is thought to be formed by colluvial slopewash processes. The upper part of the sequence contains charcoal, which could be of anthropogenic origin, and so may represent stratified archaeological sediments.

4.2.3 Archaeological Potential

4.2.3.1 Archaeological material present within the three sedimentary groups represented within this borehole is likely to have been reworked. There may, however, be locally *in situ* archaeological material on bed surfaces. The recent topsoil may also preserve features cut into the underlying units.

4.2.4 Summary

- 2.08m of Quaternary deposits were recovered.
- Three groups of sediments are present.
- A thin fluvial sand is present immediately above bedrock.
- Colluvial or solifluction sediments are present above the sands.
- Stratified archaeological material may occur at the top of the sequence.
- Archaeological potential is moderate and most material is likely to be reworked or in secondary context.




Figure 4 Stratigraphic record sheet for Borehole ARC 0006 SA

Note:
This figure shows the sample positions within the unconsolidated sediments. Samples taken from the Chalk bedrock are not shown on this figure.

4.3 ARC 0007 SA (Figure 5)

4.3.1 Stratigraphy

4.3.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.0m below ground surface. Ground level at the top of the borehole was +18.514m O.D. and the base of the borehole lay at +8.514m O.D. A total depth of c. 1.84m of unconsolidated sediments were found overlying rotted Chalk bedrock. Chalk bedrock lies at a depth of c. 1.84m (c. +16.674m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment was not clearly observed during field logging and was located immediately below bulk sample 4.

4.3.1.2 Three U4/U100 core samples were taken and 7 bulk samples recovered. Locations of samples retrieved from the unconsolidated sediments relative to the stratigraphy are shown in Figure 5. This borehole is dominated by two groups of sediments.

4.3.1.3 The first group consists of a thin silty sand unit containing chalk gravel clasts between core depths of 1.84 and 1.60m (+16.674m O.D. and +16.914m O.D.). The main group present within the borehole is seen between core depths of 1.60m and 0.34m (+16.914m O.D. and +18.174m O.D.), and consists of silty sands with sub-angular flint and chalk gravel clasts and a chalky gravel unit.

4.3.1.4 Modern roots and clay coatings are present within the upper units of the borehole between core depths of 1.16 to 0.40m (+17.354m O.D. to +18.114m O.D.). This part of the sequence may have been subjected to weathering and/or pedogenesis.

4.3.1.5 U4₁ and U4₂ both contain CaCO₃ (react with dilute hydrochloric acid), it is currently unknown whether CaCO₃ is present within the unconsolidated sediments below this point within the borehole.

4.3.2 Interpretation

4.3.2.1 The silty sandy unit containing gravel clasts (overlying the Chalk bedrock) may represent fluvial deposits. This deposit is overlain by silty sands and a gravel containing sub-angular clasts that may have been deposited by colluviation or solifluction.

4.3.3 Archaeological Potential

4.3.3.1 Any archaeological material present within the two sedimentary groups represented within this borehole is likely to have been reworked. There may be locally *in situ* material on bed surfaces. The recent topsoil may also preserve features cut into the underlying units.

4.3.4 Summary

- 1.84m of Quaternary deposits were recovered
- Two groups of sediments are present.
- A thin fluvial silty sand is present immediately above bedrock.
- Colluvial or solifluction sediments are present above the sands.
- Archaeological potential is moderate and most material will be reworked.




Figure 5 Stratigraphic record sheet for Borehole ARC 0007 SA

Note:
This figure shows the sample positions within the unconsolidated sediments. Samples taken from the Chalk bedrock are not shown on this figure.

4.4 ARC 0008 SA (Figure 6)

4.4.1 Stratigraphy

4.4.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.0m below ground surface. Ground level at the top of the borehole was +18.302m O.D. and the base of the borehole lay at +8.302m O.D. A total depth of c. 3.85m of unconsolidated sediments were found over fractured Chalk bedrock.

4.4.1.2 Five U4/U100 core samples were taken and 3 bulk samples recovered. Locations of samples retrieved from the unconsolidated sediments relative to the stratigraphy are shown in Figure 6.

4.4.1.3 Chalk bedrock lies at a depth of 3.85m (+14.452m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment is well preserved at the base of U4₅ (3.85-3.40m). The contact is sharp and smooth.

4.4.1.4 Immediately overlying the Chalk bedrock are a series of predominantly massively bedded medium to coarse sands with occasional flint gravel clasts. Towards the top of this group there are indications of cross-stratification between core depths of 2.64 and 2.38m (+15.662m O.D. and +15.922m O.D.). Clay coatings on sand grains are noted at the base of this part of the sequence and between depths of c. 3.20m and 2.30m below the top of the borehole.

4.4.1.5 The second group of sediments consists of two depositional cycles. The first commences with a gravel at the base, passes through medium to coarse sands and ends with laminated sands and silts. The second of these cycles shows the gravel to be directly overlain by the laminated silts and sands. The first of these cycles is present between core depths of 2.38m and 1.60m (+15.922m O.D. and +16.702m O.D.). The second of these cycles is composed of finer grained sand units and interbedded silts, between core depths of 1.60 and 1.00m (+16.702m O.D. and +17.302m O.D.). Sands are present between 1.00m (+17.302m O.D.) and the base of the fill at 0.28m (+18.002m O.D.). These cycles contrast with the structureless sand unit below.

4.4.1.6 Modern roots and voids (probable root canals) are present within the upper 0.30m of the borehole (+18.002m O.D. to +18.302m O.D.). Clay coatings are present below 1.00m depth and are associated with the gravel at the base of the lowest cycle.

4.4.1.7 The sediments within U4₁ (1.45-1.00m below ground surface, +16.852m to +17.302m O.D.), show a variable reaction with hydrochloric acid indicating a patchy distribution of CaCO₃. There is no reaction between the sediment and hydrochloric acid in any of the U4/U100 core samples below this depth.

4.4.2 Interpretation

4.4.2.1 The first of the sedimentary groups present within the borehole is thought to represent medium to coarse fluvial sands deposited in moderate energy conditions - probably rapidly. The second sedimentary group is formed by two sequences which show high energy conditions at the base of the sequence


depositing gravels followed by intermittent high and low energy events.

4.4.3 Archaeological Potential

4.4.3.1 The basal sedimentary group could contain reworked archaeological material or locally *in situ* material. The cyclical sequences, where gravels (with clay coatings to the clasts) are associated with the base of the cycles, may contain *in situ* archaeology associated with the surface of the gravel units. Further evidence for clay coatings in association with the upper, laminated parts of the sequence suggests further weathering. The recent topsoil may contain archaeological material.

4.4.4 Summary

- 3.85m of Quaternary deposits were recovered.
- Two groups of sediments are present.
- A massively bedded sand with gravel clasts is present immediately above bedrock.
- The upper part of the profile is dominated by cyclical sequences of interbedded gravels, sands and silts.
- Archaeological material may exist *in situ* within the bedded sands and gravels. Reworked material may exist throughout the sequence.



- 4.5 ARC 0009 SA (Figure 7)
- 4.5.1 Stratigraphy
- 4.5.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.0m below ground surface. Ground level at the top of the borehole was +16.800m O.D. and the base of the borehole lay at +6.800m O.D. A total depth of c. 4.50m of unconsolidated sediments were found overlying rotted Chalk bedrock.
- 4.5.1.2 Six U4/U100 core samples were taken and 8 bulk samples recovered. Locations of samples retrieved from the unconsolidated sediments relative to the stratigraphy are shown in Figure 7.
- 4.5.1.3 This borehole is dominated by three groups of sediments. Considerable detail exists within each group.
- 4.5.1.4 Chalk bedrock lies at a depth of 4.50m (+12.300m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment is not clearly seen and is present immediately below bulk sample 6 (B₆ on Figure 7).
- 4.5.1.5 Sediments directly overlying the Chalk bedrock consist of a series of bedded sands and gravels between core depths of 4.50 and 2.50m (+12.300m O.D. and +14.300m O.D.). The gravel at the base of the group is a sandy gravel with clasts of flint, quartz and sandstone, it is overlain by coarse sands with quartz and flint gravel clasts. Throughout this group of sediments there is a reduction in the maximum clast size present within the deposits up through the borehole to a depth of 2.50m.
- 4.5.1.6 The second of the sedimentary groups is found between core depths of 2.50m and 1.00m (+14.300m O.D. and +15.800m O.D.), and consists of laminated fine sands and silts overlying a coarse sandy gravel. A further sand overlain by bedded units occurs at the top of this part of the profile (see Plates 3 and 4). Clay coatings are found associated with the laminated parts of the sequence.
- 4.5.1.7 The third group consists of sands with gravel clasts and silty sands with sub-angular/sub-rounded gravel clasts and is present between core depths of 1.00m and 0.30m (+15.800m O.D. and +16.500m O.D.). This unit contains vertical channels that are possibly infilled root canals. Clay coatings are common throughout this part of the sequence (2.50-0.30m).
- 4.5.1.8 The borehole contains CaCO₃ (the sediment displays a patchy reaction with dilute hydrochloric acid), towards the base of U4₂ (1.45-1.00m below ground surface, +15.350 and +15.800m O.D.). The top of U4₅ (3.25-2.80m below ground surface, +13.550 and +14.000m O.D.), also reacts with hydrochloric acid. There is no reaction with hydrochloric acid within any of the other U4/U100 samples from this borehole.

4.5.2 Interpretation

4.5.2.1 The lowest group present (above the Chalk) is thought to have been deposited in a moderate to high energy fluvial environment. The reduction in the maximum clast size up through the group may indicate that the strength of the current is decreasing or the channel is shallowing. The finer laminated deposits present within the second group indicates possible floodplain/overbank channel marginal deposits. The upper group is interpreted as of colluvial or solifluction origin.

4.5.3 Archaeological Potential

4.5.3.1 The basal sedimentary group could contain reworked archaeological material. The second group may contain reworked material with the possibility of *in situ* material. The third group could again contain reworked material and *in situ* material associated with bed surfaces. The recent topsoil may also preserve archaeological features cut into the underlying units.

4.5.4 Summary

- 4.50m of Quaternary deposits were recovered.
- Three groups of sediments are present.
- Sands and gravels are present immediately above bedrock.
- The middle part of the profile is dominated by cyclical sequences of interbedded gravels, sands and silts.
- An upper group exists which may have been deposited by colluvial or solifluction processes.
- Archaeological material could exist *in situ* within the bedded sands and gravels. Reworked material may exist throughout the sequence.




Figure 7 Stratigraphic record sheet for Borehole ARC 0009 SA



Plate 4. U4 4 core at 2.20–2.65m below ground surface, borehole ARC 0009 SA showing laminated sands and silts overlying a thin gravel bed.



Plate 5. U4 3 core at 1.60–2.05m below ground surface, borehole ARC 0009 SA showing finely laminated sands and silts.

4.6 ARC 0010 SA (Figure 8)

4.6.1 Stratigraphy

4.6.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.2m below ground surface. Ground level at the top of the borehole was +18.752m O.D. A total depth of c. 2.05m of unconsolidated sediments were found overlying rotted Chalk bedrock.

4.6.1.2 One U4/U100 core sample was taken and 11 bulk samples recovered. Locations of samples retrieved from the unconsolidated sediments relative to the stratigraphy are shown in Figure 8.

4.6.1.3 This borehole contains two groups of sediments.

4.6.1.4 Chalk bedrock lies at a depth of 2.05m (+16.702m O.D.) from the top of the borehole. The contact between the Chalk and the overlying superficial sediments are sharp and were clearly seen during field logging.

4.6.1.5 Directly overlying the Chalk between core depths of 2.05m and 1.80m (+16.702m O.D. and +16.952m O.D.) is a sandy gravel with chalk and flint gravel clasts. This is overlain by the second sediment group present at depths of 1.80m to 0.26m (+16.952m O.D. and +18.492m O.D.). The group is dominated by a silty sandy with subangular to subrounded flint and chalk gravel clasts, some bedding is also present within the unit between core depths of 1.60m and 1.30m (+17.152 m O.D. and +17.452m O.D.).

4.6.1.6 Modern roots are present within the upper unit and this is seen between 0.60m and 0.40m (+18.152m O.D. and +18.352m O.D.) and there are no visible clay coatings within the profile.

4.6.1.7 The sediment shows a strong reaction with dilute hydrochloric acid within U4₁ (0.45-0.25m below ground surface, +18.302 to +18.502m O.D.). It is currently not known whether CaCO₃ is present below this depth.

4.6.2 Interpretation


4.6.2.1 The basal sandy gravel is thought to have been deposited in a moderate to high energy fluvial environment. The overlying unit is considered to be a colluvial or solifluction deposit.

4.6.3 Archaeological Potential

4.6.3.1 The stratigraphy adjacent to this borehole is likely to predominantly contain reworked archaeological material with the possibility of *in situ* material on bedding surfaces. The recent topsoil may also preserve later prehistoric archaeological features cut into the underlying units.


4.6.4 Summary

- A 2.05m deep sequence of Quaternary deposits is present.
- Two groups of sediments are present.
- A thin sand and gravel unit is present immediately above bedrock.
- Colluvial sediments are present above the basal gravel.
- Archaeological material may exist *in situ* locally. Reworked archaeological material may exist throughout the sequence.



- 4.7 **ARC 0011 SA (Figure 9)**
- 4.7.1 **Stratigraphy**
- 4.7.1.1 Ground level at the top of the borehole was +16.648m O.D. and the base of the borehole lay at +6.648m O.D. A total depth of c. 4.75m of unconsolidated sediments were found overlying rotted Chalk bedrock.
- 4.7.1.2 Nine U4/U100 core samples were taken and 7 bulk samples recovered. Sample positions within the unconsolidated sediments are shown in Figure 9.
- 4.7.1.3 This borehole contains four groups of sediments. Considerable stratigraphic detail exists within each group.
- 4.7.1.4 Chalk bedrock lies at a depth of 4.72m (+11.928m O.D.) below surface. The contact between the Chalk and the overlying sediment is poorly preserved at the base of U₄, (4.75-4.30m).
- 4.7.1.5 Immediately above the bedrock contact a poorly bedded coarse sandy gravel was present. This unit showed colour variation and contained flint, quartzite and sandstone clasts. This unit was overlain by a structureless moderately sorted medium sand with occasional gravel clasts (seen in U₄). These two units constitute the basal group of sediments. The upper surface of this group lies at a depth of 3.64m below ground surface (+13.008m O.D.).
- 4.7.1.6 Overlying the basal sands and gravels are a complex of thinly bedded gravels, sands, silts and clays between a core depth of 3.64m (+13.008m O.D.) and 1.70m (+14.948m O.D.). Trends within these deposits can be noted. Towards the top of the group, two depositional cycles (2.66 to 2.36m and 2.36 to 1.70m depth) are present, showing fining upwards trends from gravels or sands to silts/silty-clays. The fine grained upper parts of these cycles are reddened and individual grains clay coated. Trends within the sequence below 2.66m depth are less clear but a shift from gravel to bedded silts is noted between core depths of 3.64 and 3.44m. A further fine grained horizon where clay coatings are recorded coincides with traces of rooting at a core depth of c. 2.66m. A single mollusc fragment (an apex of *Pupilla muscorum*) was found at a core depth of c. 2.50m.
- 4.7.1.7 The third of the sequences is present between core depths of 1.70 and 1.10m (+14.948m O.D. and +15.548m O.D.) and exhibits a slight coarsening upwards trend from a gravelly sand to a sandy gravel.
- 4.7.1.8 The upper sequence is present between core depths of 1.10 and 0.10m (+15.548m O.D. and +16.548m O.D.) and consists of a silty sand with flint and chalk gravel clasts overlain by chalky gravels.
- 4.7.1.9 Rooting and bioturbation are present within the upper 1.70m of the borehole. Modern roots are present to depths of c. 1.00m below surface. Below 1.00m root canals are present that contain topsoil fills. Clay coatings on sand particles were noted at intervals down through the borehole (see Figure 9).

- 4.7.1.10 The sediment within the upper three U4/U100 core samples (U4₁ to U4₃ (upper portion of the U4/U100 core only)) from borehole ARC 0011 SA shows a reaction with dilute hydrochloric acid. This shows that between 1.11 and 0.10m, below ground surface (+15.538 and +16.548m O.D.) there is CaCO₃ present within the sediment. The sediment shows no reaction with dilute hydrochloric acid in any of the U4/U100 core samples below 1.11m below the ground surface (+15.538m O.D.).
- 4.7.2 Interpretation
- 4.7.2.1 It is likely that most of the sequence was deposited by moving water. The basal sediments between Chalk bedrock and a depth of 3.64m may have been deposited in a channel bar situation in moderate to high energy fluvial channel systems. The semi-cyclical sequences recorded between depths of 3.64m and 1.70m may represent deposition in a floodplain situation where repetitive shifts in channel location initiated progressive shifts to finer deposition, and subsequent exposure and weathering of the fine sediments. The third group within the sequence, between 1.70m and 1.10m, may represent a return to higher energy channel environments. The upper sequence from the borehole is dominated by coarse grained sediments containing sub-angular clasts, and may have been deposited by colluvial or solifluction processes. The recent topsoil may also preserve features cut into the underlying units.
- 4.7.2.2 Despite the presence of a single mollusc at a depth of c. 2.50m, it is unlikely that significant assemblages of vertebrate or molluscan fauna will be present in this stratigraphy.
- 4.7.3 Archaeological Potential
- 4.7.3.1 *In situ* archaeological material could occur locally in association with the surfaces of the thin gravel beds. Reworked material may occur in association with the remaining units.
- 4.7.4 Summary
- 4.75m of Quaternary deposits were present.
 - Four groups of sediments were recorded.
 - Poorly bedded sands and gravels are present immediately above bedrock.
 - Rhythmic sequences of interbedded gravels, sands and silts overlie the basal sands and gravels.
 - The upper parts of the sequence exhibit coarsening upwards trends and a flint/chalk rich gravel.
 - Archaeological material may exist *in situ* within the bedded sands and gravels. Reworked material may exist throughout the sequence.






Plate 6. Borehole ARC 0011 SA, U4 3, 22-23cm within U4/U100 core, smooth surface Quartz grain (magnification x250).



Plate 7. Borehole ARC 0011 SA, U4 4, 21-22cm within U4/U100 core, clay coated quartz grains (magnification x170).

4.8 ARC 0012 SA (Figure 10)

4.8.1 Stratigraphy

4.8.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.0m below ground surface. Ground level at the top of the borehole was +14.769m O.D. and the base of the borehole lay at +4.769m O.D. A total depth of c. 2.00m of unconsolidated sediments were found overlying rotted Chalk bedrock.

4.8.1.2 Three U4/U100 core samples were taken and six bulk samples recovered. Sample positions within the unconsolidated sediments are shown in Figure 10. Chalk bedrock lies at a core depth of c. 2.00m (c. +12.769m O.D.) from the top of the borehole. The contact between Chalk and the overlying sediment was not clearly seen but is located at the base of bulk sample 3.

4.8.1.3 The basal group of sediments is dominated by a poorly bedded sandy gravel with flint, quartz and chalk clasts, between core depths of 2.00 and 0.85m (+12.769m O.D. and +13.919m O.D.) (see Plate 8). This contains a thin sandy unit at a core depth of 1.20m. Clay coatings are present on the surfaces of the quartz grains within this sand unit (1.20 to 1.12m). The flint gravel unit at core depths of 1.12-0.85m (+13.649m O.D.), contained a small percentage of exotic clasts (i.e. non - flint/chalk/quartz/quartzites). These included sandstones and other unidentified igneous and sedimentary rock clasts.

4.8.1.4 The second sediment group is present between core depths of 0.85m and 0.35m (base of the fill) (+13.649m O.D. and +14.419m O.D.), and is a sand with subangular flint, chalk and quartz gravel clasts. Rooting (modern) is also present within the upper group, between depths of 0.80 to 0.40m.

4.8.1.5 The matrix of the gravel within U4₃ (1.60-1.45m below the ground surface, +13.169 to +13.319m O.D.) shows a reaction with dilute hydrochloric acid. There is however no reaction within the two U4/100 samples from higher within the sequence.

4.8.2 Interpretation


4.8.2.1 The basal group of sediments is thought to represent deposition in a moderate to high energy fluvial channel. The upper group is also interpreted as fluvial in origin, with the finer grain size indicating a reduction in current energy.

4.8.3 Archaeological Potential

4.8.3.1 Both of the sediment groups within the borehole may contain reworked archaeological material. However, *in situ* archaeological material may occur in association with landsurfaces (possibly shown by the clay coatings on the grains within the sandy units). The modern topsoil may also preserve later prehistoric or historic archaeological features cut into the underlying units.

4.8.4 Summary

- 2.00m of Quaternary deposits were recorded.
- Two main groups of sediments are present.
- Sands and gravels are present immediately above Chalk bedrock.
- The upper part of the profile contains a sand with sub-angular flint clasts.
- Archaeological material could exist within the sequence but is likely to be reworked. *In situ* archaeological material may occur in association with the zone of weathering/pedogenesis.



Note:
 This figure shows the sample positions within the unconsolidated sediments. Samples taken from the Chalk bedrock are not shown on this figure.

Figure 10 Stratigraphic record sheet for Borehole ARC 0012 SA



Plate 8. Unsieved bulk sample B4, ARC 0012 SA, 1.80–2.00m depth.

4.9 ARC 0013 SA (Figure 11)

4.9.1 Stratigraphy

4.9.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.0m below ground surface. Ground level at the top of the borehole was +14.159m O.D. and the base of the borehole lay at +4.159m O.D. A total depth of c. 3.30m of unconsolidated sediments were found overlying rotted Chalk bedrock.

4.9.1.2 One U4/U100 core sample was taken and five bulk samples were recovered. Sample positions relative to the unconsolidated sediments are shown in Figure 11.

4.9.1.3 Chalk bedrock lies at a depth of c. 3.30m (c. +10.859m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment was not clearly seen and is present at the base of bulk sample 3.

4.9.1.4 Immediately above the bedrock contact a poorly sorted coarse sandy gravel was present. This unit contained flint and quartzite clasts and was present between core depths of 3.30 and 0.50m (+10.859m O.D. and +13.659m O.D.). This unit also contained some exotic clasts of sandstone and other unidentified igneous and sedimentary rock clasts.

4.9.1.5 The second unit is a thin bed of silty sand with rare flint gravel clasts, present between core depths of 0.50m and the base of the modern topsoil at 0.40m (+13.659m O.D. and +13.759m O.D.). No rooting or clay coatings were noted in this sequence.

4.9.1.6 No U4/U100 samples were taken from within the unconsolidated sediments of borehole ARC 0013 SA and therefore bulk samples B₁ to B₃ were tested for CaCO₃ (reaction with dilute hydrochloric acid). B₁ and B₂ showed no reaction with dilute hydrochloric acid. B₃ (3.30 to 3.00m below ground surface, +10.859 to +11.159m O.D.) did react with hydrochloric acid. This may however be due to inclusions of the Chalk bedrock which is present immediately below the bulk sample at 3.30m below the ground surface (+10.859m O.D.).

4.9.2 Interpretation

4.9.2.1 The two sedimentary groups are thought to represent fluviially deposited sediments with some possible modification near to the surface by mass movement / re-working.


4.9.3 Archaeological Potential

4.9.3.1 The deposits could contain reworked material, and locally *in situ* archaeological material could be present in association with the surface of the gravel unit. The modern topsoil may preserve later prehistoric features cut into the underlying units.

4.9.4

Summary

- 3.30m of Quaternary deposits were recorded.
- Two groups of sediments are present.
- Sands and gravels are present immediately above bedrock.
- The upper part of the profile consists of a silty sand with occasional flint clasts.
- Archaeological material may exist within the sequence but is likely to be reworked.



Note:
 This figure shows the sample positions within the unconsolidated sediments. Samples taken from the Chalk bedrock are not shown on this figure.

Figure 11 Stratigraphic record sheet for Borehole ARC 0013 SA

4.10 ARC 0014 SA (Figure 12)

4.10.1 Stratigraphy

4.10.1.1 This borehole (see Figure 2 for location) was drilled to a total depth of 10.1m below ground surface. Ground level at the surface was +17.165m O.D. and the base of the borehole lay at +7.065m O.D. A total depth of c. 1.50m of unconsolidated sediments were found overlying rotted Chalk bedrock.

4.10.1.2 Three U4/U100 core samples were taken and four bulk samples recovered. Sample positions within the unconsolidated sediments are shown in Figure 12.

4.10.1.3 Chalk bedrock lies at a depth of 1.50m (+15.665m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment is sharp and irregular and clearly seen in the base of U4₂ (1.55-1.10m) (see Plate 8).

4.10.1.4 The stratigraphy within this borehole consists of laminated and thinly bedded fine to medium sands (with flint gravel clasts up to 2.5cm mean diameter) and silts, between core depths of 1.50m and the base of the fill at 0.50m (+15.665m O.D. and +16.665m O.D.) (see Plates 8 and 9). Sand grains are clay-coated between 0.70m and 0.50m.

4.10.1.5 There was no reaction between the sediment contained within the two U4/U100 samples taken from within the unconsolidated sediments of borehole ARC 0014 SA.

4.10.2 Interpretation

4.10.2.1 These deposits are thought to represent bedded fluvial sands and silts. Weathering and incipient pedogenesis is present at the top of the group of sediments present within the borehole.

4.10.3 Archaeological Potential

4.10.3.1 These units may contain *in situ* archaeological material. The brick rubble at the top of the borehole, 0.50-0.00m, may contain anthropogenic material from historic periods.

4.10.4 Summary

- 1.50m of Quaternary deposits were recorded.
- A single sediment unit is present.
- Laminated sands and silts are present immediately above bedrock.
- The middle part of the profile is dominated by cyclical sequences of interbedded gravels, sands and silts.
- Only the upper part of the sequence is weathered.

- Archaeological material may exist *in situ* within the bedded sands and silts. Reworked material may exist throughout the sequence. Archaeological units (associated with brick rubble) may lie immediately below the ground surface in this area.






Plate 9. U4 2 core at 1.10-1.55m below ground surface (lower part), borehole ARC 0014 SA, showing sharp contact between the sands and underlying chalk bedrock.




Plate 10. U4 2 core 1.10–1.55m below ground surface (upper part), borehole ARC 0014 SA, showing laminated sands and silts.

4.11 ARC 0015 SA (Figure 13)

4.11.1 Stratigraphy

4.11.1.1 Ground level at the top of the borehole was +12.787m O.D. and the base of the borehole lay at +2.787m O.D. A total depth of c. 2.40m of unconsolidated sediments were found overlying rotted Chalk bedrock.

4.11.1.2 Two U4/U100 core samples (of very poor quality) and six bulk samples were recovered. Sample positions relative to the unconsolidated sediments are shown in Figure 13.

4.11.1.3 Chalk bedrock lies at a depth of 2.40m (+10.387m O.D.) from the top of the borehole. The contact between the Chalk and the overlying sediment is poorly preserved at the base of bulk sample 3.

4.11.1.4 Immediately above the bedrock contact a poorly bedded coarse sandy gravel containing flint and quartzite clasts was present (see Plate 11), between core depths of 2.40 to 1.30m (+10.387m O.D. and +11.487m O.D.). A small percentage of exotic clasts of sandstones, and other sedimentary and igneous rocks were noted in this unit. This unit was overlain by a structureless moderately sorted medium sand with flint and chalk gravel clasts. A second gravel unit is present at the top of the group between core depths of 0.70 to 0.26m (+12.087m O.D. and 12.527m O.D.). Modern roots penetrate through the upper 0.70m of the stratigraphy. No clay coatings were observed on any sands within the samples.

4.11.1.5 No U4/U100 samples were successfully recovered from the unconsolidated sediments of borehole ARC 0015 SA. The 3 bulk samples recovered from the unconsolidated sediments were tested for CaCO_3 using dilute hydrochloric acid. There was no reaction with the hydrochloric acid from any of the bulk samples taken from within the unconsolidated sediments.

4.11.2 Interpretation


4.11.2.1 The sediments are thought to represent deposition in a moderate to high energy fluvial channel.

4.11.3 Archaeological Potential

4.11.3.1 Most archaeological material present within these units is likely to be reworked. Locally *in situ* finds may be present. The recent topsoil may preserve archaeological features cut into the underlying units.

4.11.4 Summary

- 2.40m of Quaternary deposits were observed.
- A single poorly bedded sandy gravel is present.
- Archaeological material is likely to be reworked throughout the sequence.



Note:
 This figure shows the sample positions within the unconsolidated sediments. Samples taken from the Chalk bedrock are not shown on this figure.

Figure 13 Stratigraphic record sheet for Borehole ARC 0015 SA



Plate 11. Unsieved bulk sample B3, ARC 0015 SA, 2.10–2.40m depth.

5 LOCATION OF TEST PITS

- 5.1 On the basis of the information obtained during the borehole phase of the evaluation it was possible to determine the location of the test-pits to be excavated in the second stage of evaluation (Figure 2).
- 5.2 It was decided that three areas required visual examination by test-pitting. Table 1 summarises the key stratigraphic elements that were expected to be located.

| | Test Pit ARC 2001 TP | Test Pit ARC 2002 TP | Test Pit ARC 2003 TP | Test Pit ARC 2004 TP |
|---|--|--|--|----------------------|
| Location | Adjacent to chalk 'cliff' between ARC 0007 SA and ARC 0008 SA | In central area of site between ARC 0009 SA and ARC 0011 SA | At western end of site between ARC 0012 SA and ARC 0013 SA | To be determined |
| Nature of sequence | Gravels and bedded sands/gravels | Bedded sands and gravels over coarse basal gravels | Coarse sandy gravels | ? |
| Depth to significant sediments | c.1.0m below ground surface | c.1.0m below ground surface | c.0.5m below ground surface | ? |
| Archaeological Potential (for Pleistocene deposits) | <i>In situ</i> material may occur in association with clay-coated gravel units | <i>In situ</i> material may occur in association with clay-coated gravel units | Predominantly reworked material may exist within the gravels | ? |

Table 1 Summary of the key stratigraphic elements that were expected to be located during test-pitting, with depths to significant Pleistocene sediments and estimated Palaeolithic archaeological potential.

- 5.1.3 The Chalk 'cliff' was broadly located between boreholes ARC0007 SA and ARC0008 SA. This is the area that had been reported in previous studies as containing the majority of the artefacts. Therefore, test pit ARC 2001 TP was excavated between boreholes ARC 0007 SA and ARC 0008 SA.
- 5.1.4 The greatest sequence complexity had been noted in boreholes ARC 0009 SA and ARC 0011 SA. Test pit ARC2002 TP was therefore excavated between these boreholes.
- 5.1.5 Boreholes ARC 0012, 0013 and 0015 SA only produced evidence for a single thick gravel accumulation. As this unit differed from that recorded in the remaining boreholes, it was decided to locate test-pit ARC2003 TP between boreholes ARC0012 SA and ARC0013 SA to investigate it.
- 5.1.6 The position of the final test-pit, ARC2004 TP, was not pre-determined. It was to be located to examine any emergent areas of interest after the excavation

of ARC 2001 - 2003 TP. In the event, it was located to the south-east of ARC2001 TP, to investigate further the line of the Chalk 'cliff'.

6 RESULTS: TEST PITS

6.1 Test-pit ARC2001TP (Figure 14)

6.1.1 This test-pit was excavated in order to confirm the nature of the stratigraphy in borehole ARC0008SA, to attempt to locate the 'cliff' line and to determine the presence or absence of archaeological deposits.

6.1.2 The test-pit was excavated with its long axis aligned from north-east to south-west and was 2.95m deep.

6.1.3 The south-west facing section was drawn (Figure 14) and the deposits recorded are detailed in Table 2.

6.1.4 Chalk bedrock was encountered at between 15.93 and 15.73m OD.

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|-------------|--|
| 0.00 - 0.25 | 100 | - | 10YR 3/2 very dark greyish brown, sandy silt with modern roots. Structureless, loose and unconsolidated. Occasional angular to sub-angular flints (<5cm). Occasional charcoal flecks and chalk clasts. Abrupt Contact |
| 0.30 - 0.50 | 101 | Spit 1 | 10YR 4/6 dark yellowish brown, silty sand to sandy silt. Loose, structureless and unconsolidated. Modern roots are present. Occasional small chalk clasts (<5cm). Diffuse Contact |
| 0.50 - 0.90 | 101 | Spit 2 | As above but hard, firm and compact. Frequent sub-angular to sub-rounded flint clasts. Modern roots. Not observed |
| 0.90 - 0.95 | 102 | Spit 3 | 7.5YR 6/8 strong brown sand. Firm and compact. No apparent structure. Occasional sub-angular to sub-rounded flint clasts. Modern roots. Diffuse contact |
| 0.95 - 1.20 | 102 | Spit 3 | As above, becoming more sandy with depth. Possibly fine bedded with alternating fine/coarse units. Modern roots still present. Sharp contact |

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|-------------|--|
| 1.20 - 1.45 | 103 | Spit 4 | 10YR 5/6 yellowish brown, mid/coarse sand. Loose and unconsolidated. In places thin bedded with 7.5YR 5/8 strong brown sand (possibly with clay coatings?). Occasional small sub-rounded flint clasts (<2cm) Sharp contact |
| 1.45 - 1.80 | 104 | Spit 5 | 7.5YR 5/8 strong brown, moderately well-sorted flint gravel. Matrix supported - matrix sandy in places varying to clay silt. Clasts <1cm to >5cm. Clasts less than 3cm are well rounded. Clasts above 3cm are sub-rounded to sub-angular. Occasional chalk clasts. Moderately compact. Interbedded with 10YR 7/6 yellow medium/coarse clean sand. Occasional mottled clay silt clasts (7.5YR 5/8) - rounded. Bedding becomes more pronounced with depth. In places sandy beds are hard, possibly indurated/cemented. Bedding alternates between gravel with sand matrix and sand with some gravel clasts. Beds dip to west. Occasional clay-silt clasts with blocky structure and open root hole traces present Sharp contact |
| 1.80 - 2.14 | 105, 106 | Spit 6 | 10YR 5/6 yellowish brown coarse sand with fine gravel. Structureless, loose and unconsolidated. Sub-angular to well rounded flint clasts (<0.5cm to +3cm). Frequent clasts of clay-silt. Becomes coarser and more poorly sorted with depth. Sharp contact |
| 2.14 - 2.30 | 107 | Spit 7 | 10YR 5/2 greyish brown, clay-silt with black flecks. Occasional rounded flint clasts (2-4cm). Frequent clay-silt inclusions (2-3cm). Beds dip to east. Thin bedded and sub-parallel. Sharp contact |
| 2.30 - 2.48 | 108 | Spit 7 | 10YR 5/2 greyish brown, sandy silt with 10YR 6/6 yellowish brown, silt. Coarse bedded - dips to west. Occasional well rounded flint clasts (<2cm). Rare sub-rounded clay-silt clasts. Diffuse west dipping contact |
| 2.48 - 2.69 | 109 | Spit 8 | 10YR 6/6 yellowish brown, medium/coarse sand with 10YR 5/8 yellowish brown staining. No apparent structure. Occasional well rounded flint clasts (<2cm). Soft, loose and unconsolidated. Diffuse west dipping contact |

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|----------------|--|
| 2.69 - 2.73 | 110, 111 | Spit 8 | 7.5YR 5/8 strong brown, medium/coarse sand. Possibly some clay silt present. Firm and compact. Frequent rounded flint clasts (<2cm). Thickens and becomes coarser with frequent gravel clasts to east (Context 110). Mixed with 10YR 5/3 brown clay-silt with some sand and clay-silt clasts. Sharp dipping contact |
| 2.73- 2.95 | 112 | Hand excavated | 10YR 5/3 brown clay-silt interbedded with 2.5Y 7/4 pale yellow sand. Very thin bedded with sub-parallel, wavy and discontinuous laminae. No clasts. Infills hollow in chalk. |
| 2.95- | | | Chalk bedrock |

Table 2 Deposits recorded in ARC2001 TP

6.1.5 Interpretation

6.1.5.1 A sequence of alternating bedded sands and silts with occasional gravel beds, represented by Contexts 112, 111, 110, 109, 108, 107, 106, 105, 104, 103, are present between 2.95 and 1.20m (15.93m and 17.86m OD). The finer deposits present indicate possible overbank, floodplain deposits. Sands and silty sands are present between 1.20 and 0.30m (17.86 and 18.58m OD) and are interpreted as colluvial deposits. The sequence of deposits is comparable to that seen in borehole ARC0009 SA.

test pit ARC 2001

K Kubiena samples

M Monolith samples




Figure 14. South-west facing section of Test Pit ARC2001 TP.

6.2 Test-pit ARC2002TP (Figure 15)

- 6.2.1 This test-pit was excavated in an area where the most complex, and deepest, stratigraphy had been recorded in an attempt to obtain the most complete profile possible.
- 6.2.2 The initial location (marked ARC2002a TP on Figure 2) of this test-pit was abandoned after excavation to a depth of 0.4m. A very mixed layer containing modern debris was encountered at this depth and, because of the danger of contamination and in consultation with a URL safety officer, it was decided to re-locate the test-pit 10m to the north-west.
- 6.2.3 The re-located test-pit was excavated with its long axis aligned from north-east to south-west and was 2.4m deep. At this depth, the side of the test pit collapsed due to the unconsolidated nature of the sediments and the test-pit was abandoned and back-filled. Below 1.2m, safety considerations precluded access and detailed recording of the stratigraphy *in situ* was not possible.
- 6.2.4 The south-west facing section was drawn (Figure 15) and the deposits recorded are detailed in Table 3.

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|------------------|--|
| 0.00 - 0.30 | 200 | - | Topsoil Sharp contact |
| 0.30 - 0.55 | 201 | Spit 1 | 10YR 3/4 dark yellowish brown medium coarse sand. Loose and soft with frequent angular to sub-rounded flint clasts (2-5cm). Occasional chalk fragments and modern roots. Occasional struck flints. Becomes darker towards base. Diffuse contact |
| 0.55 - 0.75 | 202 | Spit 1 Spit 2 | 10YR 4/4 dark yellowish brown medium sand with occasional flint/chalk clasts. Large rolled flints from base of unit. Infilled root channels present. Loose, structureless and unconsolidated. Diffuse undulating contact |
| 0.75 - 1.16 | 204 | Spit 3 | 7.5 YR 5/8 strong brown clay-silt and some sand. Firm and compact. Blocky structure. Infilled root channels. Sharp undulating contact |

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|----------------------------|---|
| 1.16 - 1.56 | 205 | Spit 4 Spit 5 | 10YR 6/8 brownish yellow soft loose sand interbedded with 7.5YR 5/8 strong brown sand possibly with clay-silt coatings. This unit is thin bedded (beds 2-3cm thick), undulating and wavy. Possibly laminated in places. Not observed |
| 1.56 - 1.61 | 205 | Spit5 | 10YR 6/8 brownish yellow very soft well sorted sand. Thin discontinuous beds of 10YR 5/8 cemented sand with occasional root traces. Occasional rare chalk fragments (2mm). Not observed |
| 1.61 - 1.90 | 205 | Spit 5 Spit 6 | 10YR 6/6 brownish yellow soft sand. Occasional rounded flint clasts. Occasional clay-silt present. Not observed |
| 1.90 - 2.40 | 205 | Spit 6 Spit 7 Spit 8 | Mid to brown grey clay silt with some sand. Possibly thin bedded or laminated in places. Occasional flint clasts. Test pit abandoned |

Table 3 Deposits recorded in ARC2002 TP

6.2.5 Interpretation

6.2.5.1 The lower part of Context 205, from 2.40-1.90m (12.26m-12.76m OD), is comparable to the upper part of the second group of sediments seen in borehole ARC0011 SA (section 4.7.1.6), possibly deposited in floodplain conditions with repetitive shifts in channel location. The rest of Context 205, from 1.90-1.16m (12.76m-13.50m OD), probably corresponds to the third group of sediments seen in ARC0011 SA, and may have been deposited in a high energy fluvial environment. The upper part of the sequence, Contexts 204, 202, and 201, from 1.16-0.30m (13.50-14.36m OD), is interpreted as being colluvial in origin.

6.2.6 In addition, a probable cut feature, Context 206, was observed in section. Its form and orientation are not known, but it may be a small ditch or gully. It was overlain by colluvial sand with occasional chalk fragments (Context 202, see Table 3). It was 0.3m deep and 0.84m wide, and contained a single fill, Context 203. This was a yellow (10YR 7/6) medium/fine sand with occasional rounded flint clasts (<5cm). A flint flake was recovered from the feature while cleaning the face of the test pit.

test pit ARC 2002




Figure 15. South-west facing section of Test Pit ARC 2002 TP.

6.3 Test-pit ARC2003TP (Figure 16)

- 6.3.1 This test-pit was excavated in an area where predominantly coarse sand and gravel deposits had been recorded in the boreholes. One of the objectives had been to establish whether the coarse sediments are bedded, which had not been possible to establish from the boreholes as only bulk samples were retrieved.
- 6.3.2 The test-pit was excavated with its long axis aligned from north-east to south-west and was 2.8m deep.
- 6.3.3 The south-west facing section was drawn (Figure 15) and the deposits recorded are detailed in Table 3.
- 6.3.4 Chalk bedrock was encountered at 11.80m OD.

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|------------------|---|
| 0.00 - 0.25 | 300 | - | Topsoil Sharp contact |
| 0.30 - 0.40/0.50 | 301 | Spit 1 | 7.5YR 5/6 strong brown medium sand. Loose and structureless with frequent poorly sorted flint clasts (<2 to +5cm), sub-angular to rounded. Common chalk clasts (<0.5 to +3cm). Modern roots present. Sharp undulating contact |
| 0.40/0.50 - 0.90 | 302 | Spit 2 | 10YR 6/6 brownish yellow medium fine sand. Soft and loose and unconsolidated. Flints still present. Chalk clasts absent. Modern roots present. Occasional struck flakes. Flint clasts increase with depth. Clasts predominantly rounded flints. Not observed |
| 0.90 - 1.54 | 303 | Spit 3 Spit 4 | 7.5 YR 5/8 strong brown to 10YR 3/6 dark yellowish brown horizontally bedded, matrix supported flint gravel. Predominantly sub-rounded to rounded flint gravel clasts (2-5cm) interbedded with structureless coarse medium sand (10YR 6/8 brownish yellow). Thin beds of clay coated sand are present and occasional quartzite clasts. Occasional modern roots. Sharp sub-horizontal contact |

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|--------------------------------------|--|
| 1.54 - 1.61 | 303 | Spit 5 | 5YR 5/8 yellowish red coarse sand with well rounded flint clasts (<1 to +5cm). Possibly clay coated sand grains. Similar to thin beds in overlying units. Sharp sub-horizontal contact |
| 1.61 - 1.75 | 304 | Spit 5 | 10YR 6/6 brownish yellow coarse to medium sand. Thin bedded with bedding dipping to west. Some 5YR 5/8 yellowish red staining present. Bedding disappears towards base of unit. Possibly becoming coarser with depth. Diffuse undulating contact |
| 1.75 - 2.81 | 305 | Spit 6 Spit 7 Spit 8 Spit 9 | Mixed 10YR 6/6 brownish yellow to 5YR 5/8 yellowish red bedded flint gravel. Poorly sorted with clasts <1cm to +10cm, rounded to angular. Occasional very large sub-angular to subrounded flint clasts (+20cm). Predominantly matrix supported. Individual beds may be present with higher sand content and clay coatings to sand grains. Firm and compact. Sharp contact |
| 2.81 - | | | Chalk bedrock |

Table 4 Deposits recorded in ARC2003 TP

6.3.5 Interpretation

6.3.5.1 Overlying the chalk bedrock in this test pit were bedded gravels and coarse sands, Contexts 305, 304, and 303, between depths of 2.81 and 0.90m (11.80m and 13.71m OD). These are similar to the basal group of sediments in borehole ARC0012 SA, and were probably deposited in a moderate to high energy fluvial environment. They were overlain by medium to fine sands, probably of colluvial origin.

test pit ARC 2003




Figure 16. South-west facing section of Test Pit ARC 2003 TP.

6.4 Test-pit ARC2004TP (Figure 17)

- 6.4.1 This test-pit was excavated in order to further investigate the area adjacent to the chalk 'cliff'. The position of the pit was determined on the basis of the deposits observed in ARC2001 TP in relation to Boreholes ARC0007 SA and ARC0008 SA.
- 6.4.2 The test-pit was excavated with its long axis aligned from north-west to south-east, at right angles to ARC2001 TP, and was 2.35m deep.
- 6.4.3 The south-west facing section was drawn (Figure 17) and the deposits recorded are detailed in Table 5.
- 6.4.4 Chalk bedrock was encountered at 16.82m OD.


| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|------------------|---|
| 0.00 - 0.30 | 400 | - | Topsoil sharp contact |
| 0.30 - 0.45 | 401 | Spit 1 | 7.5YR 4/2 brown sandy silt with occasional angular chalk fragments, angular flint clasts (<2 to +5cm). Modern roots, soft and loose. Occasional charcoal flecks and brick/tile fragments. Not observed |
| 0.45 - 0.80 | 402 | Spit 1 | 7.5YR 4/2 brown sandy silt with occasional angular chalk fragments, angular flint clasts (<2 to +5cm). Modern roots, soft and loose. Not observed |
| 0.60 - 0.80 | 403 | Spit 1 Spit 2 | 7.5YR 4/2 brown sandy silt with occasional angular chalk fragments, angular flint clasts (<2 to +5cm). Modern roots, soft and loose. Not observed |
| 0.80 - 1.00 | 405 | Spit 2 Spit 3 | 10YR 7/6 yellow silt with some sand. Becomes coarser and sandier with depth. Modern roots still present. Firm and structureless. Diffuse contact |
| 1.00 - 1.04 | 406 | Spit 3 | As above becoming 7.5YR 4/6 strong brown. Sand content decreases. Occasional black patches Sharp undulating contact |

| Depth below top of profile (metres) | Context number | Spit Number | Stratigraphic description |
|-------------------------------------|----------------|----------------------------|---|
| 1.04 - 1.80 | 407 | Spit 3 Spit 4 Spit 5 | 10YR 7/3 chalk rich sand with fine silt matrix. Matrix supported with angular to subrounded chalk clasts (<1 to +3cm). Occasional flint clasts (rounded, <2cm). Modern roots present. Becomes increasingly chalky with depth. This unit dips westwards as a body thinning to the west. Occasional flint clasts and cobbles present towards base of unit. Diffuse contact |
| 1.80 - | 409 | Spit 6 | Chalk bedrock |

Table 5 Deposits recorded in ARC2004 TP

- 6.4.5 The chalk bedrock contained pockets of sand, preserved within solution hollows. These were overlain by a chalk-rich sediment, Context 407, between depths 1.80 and 1.04m (16.82m and 17.58m OD) probably a soliflucted or colluvial deposit. It was overlain by sandy silts, 406, 405, 403, and 402, interpreted as colluvial deposits. These upper colluvial deposits are probably comparable to the main group of sediments recorded in borehole ARC0007 SA (section 4.3.1.3).
- 6.4.6 In addition, a cut feature, Context 410, was recorded at the north-west end of the test pit. It was cut from 0.80m below the modern ground surface, being overlain by colluvial sandy silt (see Table 5). Only the south-east side of the ditch, including its base, was encountered by the test pit. Judging by the profile of the side of ditch and the angle at which it was cut, it is probably 3 to 4m at the top, of which 1.7m was seen. It was 1.6m deep and contained two fills, Contexts 408 and 404. Context 408 was a strong brown (7.5YR 4/6), sandy silt with occasional angular to sub-angular flint clasts (<5 cm). Context 404 was a brown (7.5YR 4/2), sandy silt with frequent angular chalk fragments and angular flint clasts (<2 to +5cm).

test pit ARC 2004



K Kubiena samples



scale 1:20

Figure 17. South-west facing section of Test Pit ARC 2004 TP.

6.5 Summary of Artefacts and Samples Recovered

6.5.1 Table 6 summarises the depth, below modern ground surface, of the spits excavated in each test-pit, along with the deposits associated with each spit. The number of flint artefacts and pottery sherds retrieved, and bulk samples taken, is also given. Two sherds were also recovered by hand excavation from Context 404.

6.5.2 In addition to the bulk samples shown in Table 6, three monolith samples (numbers 6, 7, and 8) and four Kubierna samples (numbers 9, 10, 11, and 12) were taken from ARC2001 TP (see Figure 16), and two Kubierna samples were taken from ARC2004 TP (see Figure 19).

| Test Pit | Spit | Depth (m) | Context(s) | Flint Artefacts (no of pieces) | Pottery (No of sherds) | Bulk Sample Number |
|------------|------|-------------|---------------|--------------------------------|------------------------|----------------------|
| ARC2001 TP | 1 | 0.27 - 0.56 | 101 | 8 | 2 | - |
| | 2 | 0.56 - 0.90 | 101 | 7 | 1 | - |
| | 3 | 0.90 - 1.15 | 102 | 3 | 1 | 1 |
| | 4 | 1.15 - 1.45 | 103 | 3 | | - |
| | 5 | 1.45 - 1.70 | 104 | 2 | | 2 |
| | 6 | 1.70 - 2.11 | 105, 106 | 3 | | 3 (context 105 only) |
| | 7 | 2.11 - 2.45 | 107, 108 | 1 | | 4 |
| | 8 | 2.45 - 2.81 | 109, 110, 111 | - | | 5 (context 109 only) |
| ARC2002 TP | 1 | 0.25 - 0.55 | 201, 202 | 18 | 3 | 22 |
| | 2 | 0.55 - 0.90 | 202, 203, 204 | 1 | 1 | 23 |
| | 3 | 0.90 - 1.20 | 204, 205 | 3 | | - |
| | 4 | 1.20 - 1.65 | 205 | - | | - |
| | 5 | 1.65 - 1.90 | 205 | 1 | | - |
| | 6 | 1.90 - 2.20 | 205 | - | | 24 |
| | 7 | 2.20 - 2.40 | 205 | - | | - |
| ARC2003 TP | 1 | 0.25 - 0.55 | 301, 302 | 8 | | - |
| | 2 | 0.55 - 0.90 | 302 | 3 | | - |

| Test Pit | Spit | Depth (m) | Context(s) | Flint Artefacts (no of pieces) | Pottery (No of sherds) | Bulk Sample Number |
|------------|------|-------------|----------------|--------------------------------|------------------------|--------------------------|
| | 3 | 0.90 - 1.20 | 303 | - | | 16 |
| | 4 | 1.20 - 1.50 | 303 | 1 | | - |
| | 5 | 1.50 - 1.80 | 304, 305 | 3 | | 17 |
| | 6 | 1.80 - 2.10 | 305 | 3 | | - |
| | 7 | 2.10 - 2.35 | 305 | - | | 18 |
| | 8 | 2.35 - 2.65 | 305 | - | | - |
| | 9 | 2.65 - 2.78 | 305 | - | | - |
| ARC2004 TP | 1 | 0.30 - 0.65 | 401, 402 | 4 | | - |
| | 2 | 0.65 - 1.00 | 402 - 406 | 7 | 3 | 19 |
| | 3 | 1.00 - 1.32 | 404 - 407 | 3 | | - |
| | 4 | 1.32 - 1.75 | 404, 406 - 408 | 27 | | 20 |
| | 5 | 1.75 - 2.15 | 407 - 409 | 13 | | 21 |
| | 6 | 2.15 - 2.37 | 408, 409 | 7 | | 15 (context 408 only) |

Table 6 Summary of excavated spits, artefact recovery, and bulk samples

7 SPECIALIST REPORTS

7.1 Flint Report by M. White

7.1.1 Introduction

7.1.1.1 A collection of 130 humanly worked flints and 10 pieces of unworked burnt flint were recovered during the course of the assessment. Two general series are evident, which can be separated primarily on the basis of stratigraphic position and the presence of burnt flint, and perhaps also by condition and size. However, it is not possible to give absolute numbers for each series, as the method of recovery often cross cut natural stratigraphic units, thus obliterating the main rationale behind the division. Whilst it may have been possible to use size and condition to provide absolute counts, this was considered inappropriate without greater stratigraphic, taphonomic and numeric control. The lower series is of greatest archaeological importance and therefore forms the main concern of this report. The total collection is summarised in Table 1 and outlined in greater detail in Table 2 following the main text. In addition, a large number of small flints were retrieved during sieving of the bulk samples. However, their small size and lack of diagnostic features made it impossible to determine whether they were humanly worked, machine struck, or produced by natural processes.

| Trench | Flakes/Blades | Cores | Implements | Burnt Flint | Total |
|--------|---------------|-------|------------|-------------|-------|
| 2001TP | 27 | 1 | 0 | 2 | 30 |
| 2002TP | 22 | 1 | 0 | 6 | 29 |
| 2003TP | 16 | 2 | 0 | 0 | 18 |
| 2004TP | 54 | 3 | 4 | 2 | 63 |
| TOTALS | 119 | 7 | 4 | 10 | 140 |

Table 7 Summary of flint collection

7.1.2 Methods

7.1.2.1 Each flake was measured using dial callipers to the nearest mm. Limited technological information was also taken from all lithic components. The material recovered from the wet-sieving of the soil sample was briefly examined, but excluded as it was considered to be of minimal value.

7.1.3 Flint Series One

7.1.3.1 This collection of flints was recovered from the upper sedimentary deposits at the site, mostly from Spits 1, 2 and 3 from all trenches, and from the feature (404) cutting through trench 2004TP. It consists of small flakes, cores and one

retouched side-scraper. The collection is unstained, only mildly patinated and generally in a very fresh condition.

- 7.1.3.2 The collection exhibits mostly hard hammer flaking. There is evidence of lateral and distal flake scars on the dorsal surfaces, although the most frequent scar pattern shows parallel removals from the proximal end. Few flakes exhibit wholly cortical dorsal surfaces. The cores associated with this series show parallel, single and alternate flaking techniques (cf. Ashton 1992), totally in concord with the evidence from the flakes. The cores show between 9 and 19 flake removals each. On the basis of the small size of the flakes and cores, the relatively high number of removals per core and the high incidence of flakes with two or more distal removals, it is suggested that this series has been fairly intensively reduced from small pieces of flint, mostly using a parallel flaking technique. In addition, the cortex evident on the bulk of this series is dissimilar to that seen on flint from the Chalk in the Purfleet area, which, therefore is unlikely to have formed the source of the raw material.
- 7.1.3.3 There are no technologically or typologically diagnostic features of this series which would allow an estimate of its age, although its stratigraphic position and association with burnt flint and pottery strongly indicates a later prehistoric date.
- 7.1.4 Flint Series Two
- 7.1.4.1 This collection was recovered from the lower stratigraphic units (spits 4-7) in test pits 2001 and 2004. Given the heavily rolled condition of the material recovered from the lower stratigraphic units in test pits 2002 and 2003, it is not possible to say whether it belongs to this series, although, from its stratigraphic position, it must be Palaeolithic. A proportion of this series was found within the bedded gravels and silts of 2001TP, although by far the greatest concentration was found in the chalky slope debris (?coombe rock) in 2004TP, and from small solution hollows in the underlying chalk. The collection is mildly patinated, generally in fairly sharp condition and is stained. The colour of the staining ranges from buff to dark red-brown. The series consists of cores, flakes and two bifaces, with one possible biface throughout.
- 7.1.4.2 Two bifaces and a possible biface rough-out were recovered during the course of the assessment, all from trench 2004. The bifaces have been finished using the soft hammer knapping technique and are in fairly sharp condition. The first biface is a complete, sub-cordiform type 120mm in length. Working is restricted to the tip of the piece and the butt is mostly unworked. The cortex remaining on the artefact reveals that the original nodule was thin and narrow, and that the shape of the biface may have simply followed the form of the original blank. A second biface is represented by a broken butt fragment. The tip is missing, and the nature of the break suggests that this may have been broken during manufacture. This feature, known as end-shocking, usually results from inadequate support of the biface during manufacture. A possible biface rough-out was also recovered. This shows evidence of some attempt at thinning the piece, but the poor angles and general shape seem to have defeated the knapper. This piece is more abraded

than the other implements.

- 7.1.4.3 In addition to the bifaces themselves a small number of biface manufacturing flakes were recovered. These too are in relatively sharp condition, and are testament that the bifaces were manufactured at the site.
- 7.1.4.4 A significant assemblage of hard hammer flakes and three cores were also found. The cores show evidence of single, parallel and alternate flaking techniques. The dorsal patterns on the flakes, ranging from 1 to 5 scars, show previous removals originating from the distal, proximal and lateral positions. A number exhibit relict core edges with removals from the proximal and lateral margins. The flakes, therefore, also indicate the use of several reduction techniques, and the frequent rotation of the cores in the pursuit of good flaking angles. The flakes are often fairly large (mean length from spits 4-6, 2004TP=57mm), and exhibit a range of residual dorsal cortex from 0-100%. The latter is evidence that all stages of core reduction from initial decortication are present at the site. Some of the flakes may represent the initial hard-hammer roughing out stage of biface production.
- 7.1.5 Dating and Industrial Affinities
- 7.1.5.1 Since biface technology provides the defining characteristic of the Acheulean techno-complex (Wymer 1968), the presence of bifaces allows the confident designation of the Purfleet material to the Acheulean. Whilst this does give a general indication of the age of the site - it is definitely Lower Palaeolithic - it cannot be used to assign a specific date. Acheulean bifaces are found throughout south-eastern Britain in deposits ranging from 500 000 to 125 000 BP. Moreover, biface typology is of no value in assigning relative age. However, the nature of the industry is entirely consistent with Bridgland's proposed correlation with Oxygen Isotope Stage 9 (ca 300 000 BP).
- 7.1.5.2 Although the artefacts recovered would appear to fit Wymer's (1985) contention that a mixture of two industries - Clactonian (i.e. the flakes and cores) and Acheulean (the biface element) - are represented, there are no reasons at present for dividing the material in this manner. Core reduction of Clactonian character as present in this material is now widely recognised to be an integral part of the Acheulean (Ashton *et al* 1994, 585-589)
- 7.1.6 Raw Materials
- 7.1.6.1 The raw material used in this collection exhibits a cortex entirely consistent with that seen on flint from the Chalk at Purfleet. This is a black flint with a thin white cortex. The material appears to be good quality, with fine flaking properties, although much of the flint present in the Chalk today is badly frost shattered. Two artefacts were noted to have been made on bull-head flint which forms bands at the base of the Thanet sands and is widespread in river deposits of the Lower Thames. These factors suggest that the raw materials used are those immediately available in the area.

7.1.7 Discussion

7.1.7.1 The condition of the Lower Palaeolithic artefacts indicate that whilst not *in situ*, they have *not* moved very far from their point of origin. The use of local raw materials, and the presence of all stages of both core reduction and biface manufacture also indicates that a fairly integral assemblage is represented.

7.1.7.2 It is clear that the density of artefacts greatly diminishes away from the chalk banks of the palaeo-channel towards the centre of the channel itself. This suggests that human activity was taking place on or near the edge of the channel, and that primary knapping scatters have been swept up into a mass movement deposit and simply carried downslope. This suggests that more or less primary scatters may survive to the south of the current test trenches as well as in the margins of the fluvial deposits. The quantity of material recovered from 2004TP, in a very small volume of sediment, indicates the extreme richness of the site in the vicinity of the chalk 'cliff'. This is in line with previous observations.

| Trench | Spit | Flakes/Blades | Cores | Implements | Burnt Flint | Total |
|--------------|------|---------------|----------|------------|-------------|------------|
| 2001TP | 1 | 8 | 1 | - | 1 | 10 |
| | 2 | 7 | - | - | 1 | 8 |
| | 3 | 3 | - | - | - | 3 |
| | 4 | 3 | - | - | - | 3 |
| | 5 | 2 | - | - | - | 2 |
| | 6 | 3 | - | - | - | 3 |
| | 7 | 1 | - | - | - | 1 |
| 2002TP | 1 | 18 | - | - | 6 | 24 |
| | 2 | 1 | - | - | - | 1 |
| | 3 | 2 | 1 | - | - | 3 |
| | 4 | - | - | - | - | 0 |
| | 5 | 1 | - | - | - | 1 |
| 2003TP | 1 | 7 | 1 | - | - | 8 |
| | 2 | 2 | 1 | - | - | 3 |
| | 3 | - | - | - | - | 0 |
| | 4 | 1 | - | - | - | 1 |
| | 5 | 3 | - | - | - | 3 |
| | 6 | 3 | - | - | - | 3 |
| 2004TP | 1 | 4 | - | - | - | 4 |
| | 2 | 6 | - | 1 | 2 | 9 |
| | 3 | 3 | - | - | - | 3 |
| | 4 | 24 | 1 | 2 | 0 | 27 |
| | 5 | 11 | 2 | - | 0 | 13 |
| | 6 | 6 | - | 1 | 0 | 7 |
| TOTAL | - | 119 | 7 | 4 | 10 | 140 |

Table 8 Recovered flint collection divided according to individual spits

7.2 Pollen Analysis by R. Scaife

7.2.1 Introduction

7.2.2 Pollen analysis has been undertaken on two sub-samples from sediment monolith sections obtained from the middle-Quaternary sequence at Purfleet, Essex. The principal aims of this assessment are summarised as follows:

- To ascertain if sub-fossil pollen and spores are present in these sediments and if so, their state of preservation.
- If present, to provide preliminary information on the pollen taxonomic content and the possibility of correlating this sequence with other biostratigraphical assemblages.
- Does the site offer potential for more detailed and valid work in the future?

7.2.3 Methodology

7.2.3.1 Samples for pollen analysis were taken from the open section which had been cut back to expose a fresh face. This face was sampled in box monoliths for detailed laboratory examination and sampling. Two sub-samples were submitted by Dr M. Bates, University College, London for pollen analysis. These were from depths of 33-32cm and 14-13cm from the top of the monolith (ARC 2001 TP, monolith Sample 8 and monolith Sample 7 respectively). The positions of these samples are shown on Figure 14. However, it is noted here that monolith column 8 is the lower of the two samples.

7.2.3.2 Both sub-samples were from laminated/banded sandy silts of inorganic, minerogenic character resembling typical brickearth. Given this minerogenic character, rigorous pollen extraction procedures were required. These procedures followed those outlined by Moore *et al.* (1991) but with the addition of micromesh sieving (10 μ) for removal of the clay fraction. The samples of 7-8ml volume were decalcified with 10% HCL and deflocculated with 8% KOH. Coarse debris was removed through sieving at 150 μ and clay by micro-mesh (10 μ). Remaining silica (silt) was digested with boiling 40% hydrofluoric acid. Erdtman's acetolysis was also carried out although cellulose content was small. The concentrated pollen and spores were stained with safranin and mounted in glycerol jelly. This work was carried out in the Quaternary Environmental Change Research Centre of the Department of Geography, University of Southampton.

7.2.3.3 Pollen was found in only one of the two samples examined (32-33 cm). A preliminary pollen sum of 100 grains was identified and counted with an Olympus biological research microscope with phase contrast facility at magnifications of x400 and x1000. These data are presented in Table 9. Taxonomy follows that of Stace (1991) and Moore *et al.* (1991).

7.2.4 Results

7.2.4.1 Of the two samples examined, only that from 32-33cm (monolith 8) contained pollen which was well preserved. These data are given in table 1 below. The sample from 13-14cm (monolith 7) was almost devoid with two badly degraded saccate pollen grains (cf. *Pinus*) recorded. In the former, pollen was sparse but generally well preserved which enabled a preliminary count to be made. Overall, the pollen assemblage is not taxonomically diverse but is characterised by the dominance of temperate trees and shrubs. *Betula* (birch) and *Quercus* (oak) are dominant (33% and 31% of total pollen respectively). *Corylus avellana* (hazel) type (which might include *Myrica gale*) is also relatively important (12%). Other taxa recorded include *Fagus* (beech), *Fraxinus* (ash) and *Tilia* (lime) all of temperate woodland character. Poaceae (grasses) are the most important herb taxon. Other herbs by comparison are few with only single occurrences of *Plantago lanceolata* (plantain), *Rosacea* indet. and aquatic *Myriophyllum alternifolium* (water milfoil).

7.2.5 Discussion

7.2.5.1 Pollen has been previously found at Purfleet (Gibbard 1994). The pollen spectrum described here from monolith 8 clearly shows a temperate (interglacial) environment dominated by deciduous trees. Whilst *Quercus*, *Corylus avellana* and *Betula* are undoubtedly important, the small numbers of other thermophiles are nevertheless significant. *Fagus*, *Fraxinus* and *Tilia* are all known to be poor pollen producers and/or having limited pollen dispersion due to entomophily and in the case of *Fagus* large pollen grains. Thus, a mixed deciduous habitat is indicated. The absence of coniferous types is perhaps surprising when considered in relation to the many interglacial pollen (eg. Hoxnian) spectra from several other sites in eastern England. There is a possibility of contamination by recent (Holocene) pollen. This might, for example, come from tree root penetration or deep burrowing earthworms. Although this possibility is considered, it seems unlikely given the depth of the section, the fact that it was 'cut back' to a great degree and also the presence of pollen of the aquatic waterweed *Myriophyllum alternifolium* (albeit one grain), a species unlikely to have been present at the site during the more recent, Holocene, past. These factors concur with the interpretation of Bates that the sediments were laid down in a fluvial environment.

7.2.6 Summary and conclusions

7.2.6.1 Two pollen samples were examined of which only one produced sufficient numbers of pollen grains to enable identification and pollen counting. The pollen spectrum obtained is clearly dominated by pollen of trees of mixed deciduous woodland. This is typical of temperate interglacial character. From a single sample it is not possible to correlate this assemblage with other interglacial sequences from the lower Thames of eastern England as a whole. The possibility of contamination has been considered but it seems likely that the pollen does in fact represent sub-fossil pollen preserved in fluvially laid sediments

7.2.6.2 Given the above data, it is suggested that a more detailed palynological investigation should be considered. This should produce a valuable vegetation record from what appears to be a middle Quaternary interglacial period. This is further enhanced by the Palaeolithic archaeology which is also present in this sequence.

| | |
|----------------------------|----|
| Pinus | 1 |
| Betula | 33 |
| Quercus | 31 |
| Tilia | 1 |
| Fagus | 2 |
| Fraxinus | 1 |
| Corylus avellana type | 12 |
| Rosacea | 1 |
| Plantago lanceolata type | 1 |
| Poaceae | 16 |
| Myriophyllum alternifolium | 1 |
| Unidentified/degraded | 5 |
| cf Pteridium aquilinum | 3 |

Table 9 Pollen data obtained from 32-33cm (ARC 2001 TP, monolith 8)

7.3 Prehistoric Pottery by A. Barclay

7.3.1 Introduction

7.3.1.1 The evaluation produced a small assemblage (14 sherds) of prehistoric pottery, which includes only one featured sherd. Twelve sherds were recovered by hand excavation and two by sieving. Analysis of the fabrics indicates a broad late Bronze Age to early Iron Age (800-600 cal BC) date range for the assemblage.

Quantification

| Test pit | Context | No. of Sherds | Weight (g) | Comment |
|----------|---------|---------------|------------|----------------|
| 2001 | spit 1 | 2 | 1 | |
| | spit 2 | 1 | 6 | |
| | spit 3 | 1 | 1 | |
| 2002 | spit 1 | 3 | 7 | |
| | spit 2 | 1 | 1 | |
| 2004 | spit 2 | 3 | 33 | Shoulder sherd |
| | 404 | 2 | 14 | |
| | 408 | 1 | <1 | |
| Totals | | 14 | 64 | |

- 7.3.2 Fabrics and surface treatment
- 7.3.2.1 The majority of the sherds were manufactured from a range of flint tempered fabrics. Occasionally the flint temper has been mixed with sand. One sherd from context 404 was in a vesicular leached shell/organic and flint tempered fabric. With the exception of a fineware sherd (test pit 2002, spit 1) in a sandy fabric which is likely to be Iron Age, all of this material could be of late Bronze Age-early Iron Age date (cf. Barclay 1994; Brown 1988). At least three sherds are from fineware vessels and these have been smoothed or burnished.
- 7.3.3 Forms
- 7.3.3.1 The small assemblage consists of 13 body sherds and one featured sherd (test pit 2004, spit 2) identified as a shoulder from a probable coarseware jar.
- 7.3.4 Discussion
- 7.3.4.1 The assemblage is characterised by relatively small (mean weight 4.6g) body sherds from both coarse and fine ware vessels. The only featured sherd, a shoulder, is likely to come from an angular vessel of late Bronze Age-early Iron Age date. The overall range of fabrics is consistent with this date. Most of the material (11 sherds) came from the upper layers (colluvium) of test pits 2001-2 and 2004. Test pit 2001, spit 1 also produced a fragment of medieval/postmedieval tile indicating that at least some of this material is residual. Three sherds in flint tempered fabrics were recovered from ditch deposits 404 and 408 and it is possible that this feature is of a contemporary late Bronze Age-early Iron Age date.
- 7.4 Molluscs, Charred Plant Remains and Bones. Identifications and observations by M. Robinson
- 7.4.1 Molluscan remains were rare, although several of the samples produced specimens of *Ceciloides acicula*, a modern burrowing species. Sample 15 from Context 408, the lower fill of the later prehistoric feature in ARC2004 TP, produced a range of open country molluscs (*Cepaea* sp., *Vallonia excentrica*, *V. costata*, and *Pupilla muscorum*), one of shaded habitat (*Discus rotundatus*), and a single example of *Cepaea nemoralis*, a partially aquatic species. This range of species would not be incompatible with the environment pertaining in, for example, a deep, open ditch in unwooded countryside. A very small quantity of animal bone was also present, representing two species - mouse (*Apedemus* sp.) and an amphibian. The same context also contained two glume bases of spelt wheat (*Tritium spelta*), two indeterminate cereal grains, and a small quantity of oak charcoal (*Quereus* sp.).

8 SUMMARY OF RESULTS

8.1 General Stratigraphic Sequence

8.1.1 The evaluation indicates that the stratigraphic sequences observed fall into three separate spatial groups (Figure 18). Group 1 consists of the deposits observed in boreholes ARC0012 SA, ARC0013 SA, and ARC0015 SA and in test pit ARC2003 TP. These boreholes were characterised by coarse sands and gravels, likely to have been deposited under medium to high energy environments of deposition. The sediments occur between 14.50m and 10.39m OD. The sedimentology indicates that they were deposited in fluvial conditions.

8.1.2 Group 2 consists of the deposits recorded in boreholes ARC0008 SA, ARC0009 SA, ARC0011 SA, and ARC0014 SA and test pits ARC2001 TP and ARC2002 TP. A sequence of coarse fluvial sands and gravels was overlain by well bedded sands and silts with occasional thin gravel beds. These were interpreted as fluvial deposits grading vertically into channel margin or overbank floodplain deposits. There was some indication of episodes of pedogenesis and therefore of potential palaeo-landsurfaces. The upper part of the sequences in ARC0009 SA and ARC 0011 SA consisted of chalky silt or gravel. These are probably colluvial or soliflucted sediments.

8.1.3 Group 3 consists of the deposits recorded in boreholes ARC0006 SA, ARC0007 SA, and ARC0010 SA, at the eastern end of the site. Test pit ARC2004 TP can probably also be included in this group. Chalk bedrock overlaid a very thin bed of sand or gravel. This was overlain by a chalk-rich sediment, interpreted as colluvial or solifluction deposits.

8.2 Test Pits

8.2.1 ARC20001 TP did not locate the chalk 'cliff'. However, the rockhead was 1.5m higher than in the adjacent borehole ARC0008 SA (see Figure 19). The sediments present were well bedded and would be suitable for palaeo-current determinations. Flint artefacts were recovered from all the spits except Spit 7. Several flint flakes were present in the well bedded, pleistocene deposits in the lower part of the sequence. Their relatively sharp condition would indicate that they have not moved far from their point of origin. The upper deposits (Contexts 101 and 102) produced the majority of the flint artefacts, although this material is undiagnostic, it differs in character from the material in the lower levels of the sequence, and two small sherds pottery, recovered from Spits 1 and 2, indicate a late prehistoric date. The depth of Context 102 suggests that it may fill a cut feature, perhaps that seen in ARC20004 TP (Context 410), although it was not possible to be certain of this within the confines of the test pit.

8.2.2 It was difficult to record the deposits in ARC2002 TP due to the unconsolidated nature of the sediments and the depth of the sequence was not ascertained. However, the lower part of the recorded sequence (Context 205) seemed to confirm the nature of the sequence as recorded in ARC0011 SA. A single flint flake was recovered from the lower sequence (from Spit 5) but




Figure 18. Summarised stratigraphy along route corridor based on selected boreholes. N.B. An additional single borehole (RC 3506) drilled by URL as part of a previous investigation is included on this transect.

was in a rolled condition and undiagnostic. The upper deposits (Spits 1 to 3) produced a greater number of flint artefacts and a few small sherds of pottery. One flint flake was retrieved directly from Context 203, which fills what is probably a late prehistoric feature. It was cut into a firm silty clay with a blocky structure, Context 204, which may represent the contemporary ground surface. The overlying deposits, Contexts 201 and 202, are later, possibly colluvial deposits.

8.2.3 ARC2003 TP demonstrated that the coarse sands and gravels recorded in this area are in fact well-bedded and are suitable for palaeo-current determinations. A small number (7) of flint flakes were recovered from Spits 4, 5 and 6, all in a rolled condition. The upper part of the sequence (Spits 1 and 2) also produced a small quantity of flint, including two cores, which can probably be assigned to the late prehistoric period. Again, the later deposits may be colluvial in origin.

8.2.4 ARC2004 TP, immediately to the south-east of ARC20001 TP, revealed a chalky basal sediment (Context 407), possibly a colluvial or solifluction deposit. The lower part of 407 appeared to consist of mixed chalky rubble with pockets of sandy material filling hollows in the chalk bedrock. The majority of the Lower Palaeolithic artefacts came from this deposit and includes two bifaces, a possible biface rough-out, and a number of biface manufacturing flakes. They are generally in fairly sharp condition, suggesting that they have not moved far from their point of origin. The Pleistocene deposits were truncated by a large feature (Context 410) filled with brown sandy silt (Contexts 404 and 408). 404 produced two sherds of pottery of Late Bronze Age or Early Iron Age date. This feature may represent a substantial ditch but the small size of the test-pit precludes a more precise interpretation.

8.3 Biological and Artefactual Material

8.3.1 The assessment of the bulk soil samples produced little environmental material, except from the later prehistoric feature in ARC2004 TP which produced a small assemblage of molluscs, charred plant remains, small rodent and amphibian bones, and a fragment of pottery.

8.3.2 Examination of the sieved fraction of the non-floating residues of the bulk samples produced quantities of small flint flakes. These occurred in all of the samples except Sample 5 from Context 109.

8.3.3 Two of the column samples from the lower part of the sequence in ARC2001 TP (see Figure 14) were assessed by Dr R Scaife for pollen preservation. One of these, from the lowest, laminated unit (at 15.81m OD) produced a pollen spectrum dominated by pollen of trees of mixed deciduous woodland.




Figure 19. Summarised stratigraphy across southern edge of palaeochannel.

from very limited, relatively haphazard sample observations.

10.1.2 A further important issue, which to some extent overcomes the limits of the concept of archaeological remains as 'monuments' is that modern attitudes to the investigation of palaeolithic archaeology are fully integrated into multidisciplinary studies of pleistocene sedimentary and environmental sequences. As such the approach overlaps considerably with the geological and scientific interest of deposits whose importance (as in this case) may be recognised through the designation of SSSI's under a different body of legislation. It may also be noted that the Scheduled Monument criteria were based on the concepts already established for the designation of SSSI's, and so while they are different, the two approaches do at least provide a parallel framework for assessing the scientific importance of an area of interest.

10.2 The Purfleet Pleistocene and Palaeolithic Complex

10.2.1 It is understood that for the designation of the SSSI at Purfleet the area east of the Esso Pit was excluded from the SSSI designation because it was not certain at the time whether or not it was made ground (D Bridgland pers comm.).

10.2.2 Despite the difficulties of using the non-statutory scheduled monument criteria for this type of site explained above, there is a need to judge the significance of the area affected by the CTRL within the context of the whole Pleistocene deposit at Purfleet. For the purposes of this assessment OAU has used the criteria at two levels; first for the whole of the Purfleet pleistocene gravel and chalk head deposits east of the present Mardyke through which the CTRL will pass; and secondly for the specific CTRL corridor investigated for this Task. This is set out in Table 10.

10.2.3 It is concluded that the CTRL corridor is an area which forms an integral part of a complex of national importance for its pleistocene and palaeolithic interest. Within the length of corridor examined it seems clear that the areas of greatest potential are the general area of the chalk 'cliff' and in particular the solifluction deposits which are most productive of flint artifacts, the finer fluvial sediments which have greatest potential for biological remains, and the areas in the fluvial sands and gravels with signs of incipient pedogenesis which are among the sedimentary contexts which might produce *in situ* traces of palaeolithic activity. While actual *in situ* remains have not been identified here or elsewhere within the deposits at Purfleet, the freshness of the flintwork and the presence of potentially suitable sedimentary contexts leaves this as a possibility.

9 CONCLUSIONS

9.1 Pleistocene

- 9.1.1 The evaluation has produced a transect through the fluvial deposits and feather edge of the Corbets Tey Gravels, and has identified an artefact-rich, slope wash or solifluction deposit close to the line of the chalk 'cliff'. Unfortunately, it was not possible to relate this deposit to the fluvial deposits within the channel due to the presence of a large, later prehistoric feature.
- 9.1.2 The laminated and bedded silts, sand and clays seen in borehole group 1 and in ARC2001 TP have similarities with deposits recorded in Bluelands and Greenlands Quarries, where they produced mollusca, ostracods and pollen, and in the Botany and Esso Pits, where, like this sequence, the deposits were without fossils. The accepted model development of the Thames fluvial sequences (Bridgland 1994, 17-19) indicates that these deposits were laid down in interglacial, floodplain conditions. However, a discrepancy exists in that this sequence has previously been recorded at a height of about 10.6m OD, compared to between about 14.5m and 17.5m OD in the assessment area. However, since the silts, sands and clays observed in the nearby exposures are overlain unconformably by sands and gravels, it is possible that the deposits recorded during the present work represent a later part of the same interglacial sequence which has been eroded elsewhere.
- 9.1.3 Samples from the Pleistocene deposits did not produce faunal remains. However, the pollen spectrum recovered from the lower, interbedded silts in ARC2001 TP shows a temperate (interglacial) environment, dominated by trees of mixed deciduous woodland, consistent with the environment of deposition suggested by Bridgland (*op cit*).
- 9.1.4 The dating of interglacial deposits at Purfleet is uncertain - Bridgland (1994, 225-228) correlates this interglacial with Oxygen Isotope Stage 9, suggesting an age of about 300 000 BP (mid-Saalian Stage). However, Gibbard (1994) suggests that the interglacial sediments represent part of the later, Ipswichian stage. In the absence of molluscan material suitable for amino acid geochronological determinations, other dating techniques such as Electron Spin Resonance, Thermoluminescence, and Chlorine-36 techniques would need to be considered.
- 9.1.5 The form of the chalk 'cliff' was not clearly revealed, but the fluvial sediments thin out against rising chalk bedrock in the south-eastern part of the site, suggesting that the 'cliff' is fairly low relief in character and complicated by solifluction or slope wash deposits (see Figure 19).
- 9.1.6 Archaeological material was recovered from all of the test pits but that recovered from the lower parts of ARC2002 TP and ARC 2003 TP was heavily rolled and re-worked. While not specifically diagnostic as palaeolithic material, the stratigraphic position makes this likely. In ARC2001 TP, a number of flint flakes, in fairly sharp condition, were recovered from the lower silt units. In ARC2004 TP, the majority of the flint was recovered from the chalk-rich deposit overlying chalk bedrock, and from hollows within the

bedrock, again in good condition. This deposit is probably a solifluction or slope wash deposit, eroded from the chalk 'cliff'. The condition of the artefacts suggests that they have not moved very far, indicating that *in situ* deposits may exist locally, either on the margin of the floodplain deposits, or to the south of the chalk 'cliff'. This association of archaeological material with the 'cliff' line is in accord with the situation recorded in other exposures at Purfleet. For example, archaeological material was recovered from sands and gravels banked against the chalk 'cliff' at Botany Pit (Wymer, 1968, 312-313 and 1985, 313) - some of this material was associated with chalky solifluction deposits.

9.1.7 The worked flint from ARC2004 TP is considered to belong to the Acheulian techno-complex, the most characteristic form of which is the bifacial hand-axe. Two bifaces, a possible rough-out, and a small number of biface manufacturing flakes were recovered from the chalk-rich deposit in ARC2004 TP (see Plate 12). In addition, there were several cores and hard hammer flakes, of Clactonian character. This accords with material recovered from Bluelands and Greenlands Quarries from the basal gravel underlying the laminated silts, and from a gravel band overlying it. The possibility therefore exists that more than one Palaeolithic industry is represented by this material, as suggested by Wymer (1985). No material attributable to the Levalloisian industry was recovered during the present work.

9.1.8 This evaluation has confirmed that significant quantities of palaeolithic flint artefacts occur within the Pleistocene deposits affected by the CTRL at Purfleet. Their fresh condition, together with the evidence of flintworking and the range of implements and waste material present, reinforces the significance of the assemblage. Although there is no direct evidence for *in situ* from test pits 2001 and 2004, it has only travelled a short distance and the possibility of *in situ* deposits cannot be ruled out. The pollen confirms that there is at least some potential for palaeo-environmental reconstruction which can be coupled with the sedimentary analyses likely to shed light on the depositional conditions. The absence of molluscs and small bones and the very poor preservation of one of the pollen samples indicates that there is unlikely to be a continuous sequence of palaeoenvironmental evidence of all types. However, the absence of molluscs and bones in these test pits does not mean that there will not be pockets or more extensive localised sequences of rich deposits within the CTRL corridor in the vicinity of the chalk cliff.

9.1.9 The possibility that the basal sequence in ARC2001 TP may represent temperate interglacial deposits of later date than observed elsewhere at Purfleet gives this particular area a potential added significance which is not applicable to other parts of the Purfleet complex as currently understood. However, further investigations would be needed to confirm this.

9.2 Later Prehistoric

9.2.1 Although the investigation was specifically designed to assess the Pleistocene and associated Palaeolithic deposits, two later prehistoric features were recorded, in ARC2002 TP and ARC2004 TP. The large, ditch-like, feature in ARC2004 TP produced pottery of Late Bronze Age or Early Iron Age date and



Plate 12. Bifacial hand-axes (top) and biface manufacturing flakes (bottom) from ARC 2004 TP.

both features produced undiagnostic flint flakes. In both cases, the features were overlain by brown sand, or sandy silt, from 0.30 to 0.55m thick, underlying the topsoil. Similar deposits were also recorded in ARC2001 TP and ARC2003 TP. The spits corresponding to these deposits in each case produced a noticeable concentration of flint flakes, as well as a small quantity of pottery. These upper deposits have been interpreted as colluvial deposits and may correspond with similar deposits recorded at Greenlands Quarry (Gibbard, 1994, 74).

- 9.2.2 It is not known what the large ditch-like feature in ARC2004 TP may have been part of or how significant it may have been. However, it is a relatively substantial feature for this period, and compares closely with, for example, the size of the ditches of the North and South rings at Mucking (Clarke, 1993) and is much larger than, for example, the ditches of the enclosed settlement at Lofts Farm (Brown, 1988), though not dissimilar to linear later Bronze Age ditches at Gravesend (Mudd, 1994). If it were to belong to a large enclosure of the Mucking type, which it also parallels in topographical location on top of a low chalk hill overlooking the Thames, it would be of at least regional significance. Other examples of substantial late Bronze Age enclosures of this type occur at Springfield Lyons, Essex; Mill Hill, Deal and Highstead, Kent; and St Mary's Hospital, Carshalton, Surrey.
- 9.2.3 If in fact, it were later in date, the ditch might represent a smaller settlement enclosure of the more common Iron Age type, or might just be a large boundary ditch. The other small feature (Context 206) would be typical of a variety of gullies and ditches on later prehistoric sites, ranging from parts of whole field systems as at Mucking to minor lengths of gully of no obvious function.

10 SIGNIFICANCE OF THE DEPOSITS

10.1 General Considerations

- 10.1.1 The significance of archaeological remains in Britain are normally judged by reference to the Secretary of State's non-statutory criteria for the scheduling of Ancient Monuments (eg as set out in *DoE Planning Policy Guidance Note 16*) and for the purposes of their Monuments Protection Programme English Heritage have added two additional criteria (amenity and conservation value) which are used for the purpose of considering management issues. There is no standard guidance for how the criteria should be applied to 'open' palaeolithic sites (ie natural sedimentary deposits containing palaeolithic artifacts rather than more specific confined sites such as caves and rock shelters). These open sites do not readily fit within the definition of an ancient monument to which the criteria apply: although the criteria provide a relatively universal set of considerations against which the significance of archaeological remains may be judged, there is a particular difficulty with palaeolithic open sites of this type in defining the spatial limits of the area to which they are being applied. This is particularly difficult when the archaeological evidence is not necessarily co-terminus with the extent of the natural deposits in which it occurs, and where its extent can only be judged

Table 10 Assessment of the Significance of the Deposits

| Criteria | Whole Deposit | Evaluation Area |
|--------------------------|--|--|
| Survival/ Condition | Significant parts of the deposit have been quarried away in the past leaving about 27% intact (cf URL 1994, Vol 4 Figure B4). The sedimentary sequence survives well where not disturbed. In places there is a good range of biological palaeoenvironmental evidence, but its survival appears to be patchy. By virtue of the depth of the deposits they are relatively stable. In the vicinity of open faces there could be some deterioration of preserved pollen, though this has not been documented. Palaeolithic artifacts within the deposit are often very fresh though no <i>in situ</i> working floors or occupation areas have yet been identified. | The survival of pleistocene deposits within the CTRL corridor is similar to the general situation, except that the area of greatest interest for palaeolithic archaeology along the gravel/ chalk interface is apparently largely undisturbed, except for the limited effects of a probably later prehistoric ditch. |
| Period | Clactonian, Acheulian and Levalloisian artifacts reflecting the chronological development of flint working technology have been found in stratigraphic sequence, but dating deposits on the basis of artifacts relies partly on the absence of technologically later types, which may not be significant. On the basis of amino acid dating of molluscan remains and correlation of geological stratigraphies within the region the sequence is considered to belong to Oxygen Isotope Stages 10-9. There is no absolute dating for the sequence. | The presence of Acheulian type flintwork is consistent with the general period of the deposits. There are some differences in the height of the deposits which make precise correlation with the stratigraphy elsewhere difficult though the sequence is generally consistent with previous observations. Direct dating of the sequence may be difficult. |
| Rarity | Pleistocene deposits with a known abundance of fresh artifacts, demonstrable palaeoenvironmental evidence and possibly a stratigraphic sequence of technological types are very rare on a national scale. In this respect the complex is also very rare within the Thames sequence and in particular is an unusually rich and complex sequence for the particular stages of the Thames valley development represented here. | As for the general deposit with regard to the presence of fresh artifacts; the deposits investigated appear to have less potential for palaeoenvironmental evidence and stratigraphic sequencing which would diminish their rarity value if generally applicable to the deposits affected by the CTRL, but pockets of good preservation could well exist which were not encountered by this investigation. |
| Fragility/ Vulnerability | Apart from the CTRL there are some pressures from development in the area, though it is not designated for further mineral extraction. The designation of part of the area as an SSSI provides some protection. | Significant vulnerability to CTRL proposals as the railway will be in cutting through the full depth of the deposits. |

| Criteria | Whole Deposit | Evaluation Area |
|---------------|---|--|
| Diversity | The deposits incorporate a variety of depositional environments and ancient topographical features no longer visible on the surface. In places there is a good range of other (biological) palaeoenvironmental evidence. Palaeolithic artifacts have been found in significant numbers in several widely separated parts of the deposit and at different levels within it. A range of later material has come from the area, though it is not clear what surviving <i>in situ</i> archaeology might be present. | As for the general deposit, but the stratigraphic sequence appears to be somewhat less diverse and the absence of molluscan and faunal remains suggests that the evidence from this particular area may be somewhat less diverse. However pockets of better preservation could survive which were not encountered in the limited sample of the area investigated. |
| Documentation | The pleistocene sequence at Purfleet has received much attention in recent years and with the results from this investigation is one of the best documented palaeolithic sites of this type in Britain. | This report provides the bulk of the documentation for the CTRL corridor though finds from the immediately adjacent areas such as the Esso Pit are also relevant. |
| Group Value | The Purfleet deposits have significant group value because of their key position in relation to other deposits of major significance in the Lower Thames valley such as Ebbsfleet, and how the comparative geological stratigraphy and archaeology between such sites is crucial to understanding the development of pleistocene geomorphology and palaeolithic archaeology in the region (Bridgland 1994) | The deposits within the CTRL corridor have substantial group value as part of the general Purfleet deposits, and thereby as a contribution to wider understanding of the development of the Thames sequence and early human activity in the region. They have specific group value in that there are differences in the height of deposits which cannot be accounted for on the basis of the results of this investigation but could be of wider significance in interpreting the sequence as a whole. |
| Potential | The Purfleet deposits have very significant archaeological potential. It is only in recent years as a result of small scale systematic observation that the deposits are beginning to be understood. There is potential for the survival of <i>in situ</i> palaeolithic archaeology and for a significant range of good sedimentary and biological palaeoenvironmental evidence which has not yet been fully exploited. There is significant potential for testing and greatly refining present understanding of the spatial development and patterning of the ancient topography and associated palaeolithic activity which could have much wider implications for palaeolithic archaeology. There is significant potential for clarifying the human use of the site in the Holocene which at present is very poorly understood but could be of significance relative to more thoroughly investigated areas in this part of the Thames valley. | As for the general deposit and as above regarding the height of deposits within the CTRL corridor compared to other parts of the Purfleet sequence. The potential for good biological evidence as revealed by the investigation reported here is more limited than elsewhere in the Purfleet deposits, but patches of better preservation might exist. |

| Criteria | Whole Deposit | Evaluation Area |
|--------------------|--|--|
| Amenity Value | The site has no public amenity value, | As for the deposits in general. |
| Conservation Value | The Purfleet sequence has significant conservation value in relation to geological and geomorphological deposits, but is not of significant ecological interest. | As for the deposits in general. |
| Overall Assessment | The Purfleet pleistocene and palaeolithic complex is of clear national importance, as widely recognised by both geomorphologists and archaeologists with specialist expertise. The multi-disciplinary interest of the complex reinforces its significance. | The CTRL corridor traverses an integral part of the overall Purfleet complex, which has now been shown not to be significantly disturbed as previously suspected. While the potential interest of this area would be somewhat diminished if the apparent lack of molluscan and faunal remains proved to be general, the possibility of such remains being well preserved in pockets of suitable deposits cannot be excluded. The potential interest of the area is enhanced by the possible existence of a later prehistoric site of some significance. The CTRL corridor is part of a complex of national importance. |

11 BIBLIOGRAPHY

- Ashton N., Irving B., Lewis S., McNabb J., Parfitt S. 1994 'The contemporaneity of Clactonian and Acheulean flint industries at Barnham, Suffolk' *Antiquity* 68, 585-589
- Bridgland D. R. 1994 *Quaternary of the Thames* (London)
- Brown, N. 1988 'A Late Bronze Age enclosure at Lofts Farm, Essex' *Preh. Proc. Soc.* 54, 249-302
- Caldwell A. 1971 'Beacon Hill, Purfleet' *Panorama* 14, 58
- Clarke A. 1993 *Excavations at Mucking* EH Archaeological Report No 20
- Gibbard P. G. 1994 *Pleistocene History of the Lower Thames Valley* (Cambridge)
- Gilman P. J. 1956 'Purfleet: Botany' *Panorama* 2, 12-14
- Hollin J. T. 1977 'Thames interglacial sites, Ipswichian sea levels and Antarctic ice surges' *Boreas* 6, 33-52
- Moore P. D., Webb J. A. and Collinson M. E. 1991 *Pollen analysis* (Oxford)
- Mudd A. 1994 'The excavation of a later Bronze Age Site at Coldharbour Road, Gravesend' *Archaeologia Cantiana* 114, 363-410
- Union Railways Limited. 1994 *Assessment of Historic and Cultural Effects of the Channel Tunnel Rail Link*. Prepared by the Oxford Archaeological Unit
- Union Railways Limited. 1995a *Purfleet, Thurrock, Essex - Method Statement*. Prepared by the Oxford Archaeological Unit
- Union Railways Limited. 1995b *Purfleet - Site Safety Plan and COSHH Statement*. Prepared by the Oxford Archaeological Unit
- Union Railways Limited. 1995c *Purfleet, Thurrock Essex - Written Scheme of Investigation*. Prepared by the Oxford Archaeological Unit
- Palmer S. 1975 'A Palaeolithic Site at North Road Purfleet Essex' *Transactions of the Essex Archaeological Society* 7, 1-13
- Stace C. *New flora of the British Isles* (Cambridge)
- Wymer J. J. 1968 *Lower Palaeolithic Archaeology in Britain, as represented by the Thames Valley* (London)
- Wymer J. J. 1985 *The Palaeolithic Sites of East Anglia* (Norwich)

APPENDIX: Sample Statistics for Boreholes ARC 0006 SA to ARC 0015 SA

Borehole: ARC 0006 SA

Date drilled: 28-31/07/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|----------------|
| 0.40-0.65 | U4 ₁ | |
| 0.65-0.80 | | B ₁ |
| 1.05-1.25 | | B ₂ |
| 2.00-2.10 | | B ₃ |
| 2.10-2.40 | | B ₄ |
| 3.50-3.80 | | B ₅ |
| 5.10-5.60 | | B ₆ |
| 7.30-7.60 | | B ₇ |
| 9.40-9.80 | | B ₈ |

Borehole: ARC 0007 SA

Date drilled: 31/07/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|----------------|
| 0.40-0.75 | U4 ₁ | |
| 0.75-0.90 | | B ₁ |
| 0.90-1.15 | U4 ₂ | |
| 1.20-1.35 | | B ₃ |
| 1.60-1.93 | | B ₄ |
| 1.90-2.35 | U4 ₅ | |
| 2.35-2.50 | | B ₅ |
| 3.20-3.50 | | B ₆ |
| 5.40-5.70 | | B ₇ |
| 8.30-8.70 | | B ₈ |

Borehole: ARC 0008 SA

Date drilled: 24/07/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|----------------|
| 0.50-1.00 | | B ₁ |
| 1.00 - 1.45 | U4 ₁ | |
| 1.60 - 2.05 | U4 ₂ | |
| 2.05 - 2.20 | | B ₂ |
| 2.20 - 2.65 | U4 ₃ | |
| 2.80 - 3.25 | U4 ₄ | |
| 3.25 - 3.40 | | B ₃ |
| 3.40 - 3.85 | U4 ₅ | |

Borehole: ARC 0009 SA

Date drilled: 26/07/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|-----------------------------|-----------------|----------------|
| 0.40 - 0.85 | U4 ₁ | |
| 0.85 - 1.00 | | B ₁ |
| 1.00 - 1.45 | U4 ₂ | |
| 1.45 - 1.60 | | B ₂ |
| 1.60 - 2.05 U4 ₃ | | |
| 2.05 - 2.20 | | B ₃ |
| 2.20 - 2.65 | U4 ₄ | |
| 2.65 - 2.80 | | B ₄ |
| 2.80 - 3.25 | U4 ₅ | |
| 3.40 - 3.85 | U4 ₆ | |
| 4.00 - 4.45 | | B ₆ |
| 4.50 - 5.00 | | B ₇ |
| 6.70 - 6.90 | | B ₈ |
| 8.10 - 8.70 | | B ₉ |

Borehole: ARC 0010 SA
 Date drilled: 01/08/95
 Maximum depth: 10.20m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|-----------------|
| 0.25-0.45 | U4 ₁ | |
| 0.45-0.60 | | B ₁ |
| 0.60-0.80 | | B ₂ |
| 0.80-1.00 | | B ₃ |
| 1.00-1.20 | | B ₄ |
| 1.20-1.30 | | B ₅ |
| 1.30-1.50 | | B ₆ |
| 1.60-1.80 | | B ₇ |
| 1.80-2.05 | | B ₈ |
| 2.30-2.50 | | B ₉ |
| 5.30-5.60 | | B ₁₀ |
| 8.10-8.40 | | B ₁₁ |

Borehole: ARC 0011 SA

Date drilled: 28/07/95
 Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|-----------------|
| 0.10-0.30 | U4 ₁ | |
| 0.30-0.45 | B ₁ | |
| 0.45-0.90 | U4 ₂ | |
| 1.05-1.50 | U4 ₃ | |
| 1.50-1.60 | | B ₃ |
| 1.60-2.05 | U4 ₄ | |
| 2.05-2.20 | | B ₄ |
| 2.20-2.55 | U4 ₅ | |
| 2.55-2.70 | | B ₅ |
| 2.70-3.15 | U4 ₆ | |
| 3.15-3.30 | | B ₆ |
| 3.30-3.60 | U4 ₇ | |
| 3.70-4.15 | U4 ₈ | |
| 4.30-4.75 | U4 ₉ | |
| 4.75-4.90 | | B ₉ |
| 7.00-7.10 | | B ₁₀ |

Borehole: ARC 0012 SA

Date drilled: 02/08/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|----------------|
| 0.40-0.85 | U4 ₁ | |
| 0.85-1.00 | | B ₁ |
| 1.00-1.30 | U4 ₂ | |
| 1.30-1.45 | | B ₂ |
| 1.45-1.60 | U4 ₃ | |
| 1.80-2.00 | | B ₄ |
| 2.00-2.50 | | B ₅ |
| 4.90-5.20 | | B ₆ |
| 7.10-7.50 | | B ₇ |

Borehole: ARC 0013 SA

Date drilled: 24-25/07/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|----------------|
| 0.50 - 0.85 | | B ₁ |
| 1.45 - 1.70 | | B ₂ |
| 3.00 - 3.30 | | B ₃ |
| 3.30 - 4.00 | | B ₄ |
| 5.50 - 5.70 | | B ₅ |
| 5.70 - 6.25 | U4 ₁ | |

Borehole: ARC 0014 SA

Date drilled: 01/08/95

Maximum depth: 10.10m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|----------------|
| 0.50-0.95 | U4 ₁ | |
| 0.95-1.10 | | B ₁ |
| 1.10-1.55 | U4 ₂ | |
| 1.55-1.70 | | B ₂ |
| 1.90-2.35 | U4 ₃ | |
| 4.00-4.30 | | B ₃ |
| 7.10-7.30 | | B ₄ |

Borehole: ARC 0015 SA

Date drilled: 25/07/95

Maximum depth: 10.00m

| Borehole depth (m) | U4/U100 samples | Bulk samples |
|--------------------|-----------------|--|
| 0.80 - 1.00 | | B ₁ |
| 1.40 - 1.70 | | B ₂ |
| 2.10 - 2.40 | | B ₃ |
| 2.70 - 2.90 | | B ₄ |
| 3.50 - 3.95 | | U4 failed - saved as a bulk sample |
| 4.20 - 4.65 | U4 ₃ | |
| 5.50 - 5.95 | U4 ₄ | |
| 8.00 - 8.20 | | B ₅ |



OXFORD ARCHAEOLOGICAL UNIT

46 Hythe Bridge Street, Oxford, OX1 2EP

Head Office Tel: 01865 243888 Fax: 01865 793496

Post-Excavation Tel: 01865 204642 Fax: 01865 204637



Director: David Miles B.A., F.S.A., M.I.F.A.

The Oxford Archaeological Unit Limited. Registered Charity Number: 285627
Private Limited Company Number: 1618597 Registered Office: 46 Hythe Bridge Street